



UHI Research Database pdf download summary

Linking environmental variables with regional-scale variability in ecological structure and standing stock of carbon within UK kelp forests

Smale, Dan A.; Burrows, Michael T.; Evans, Ally J.; King, Nathan; Sayer, Martin D. J.; Yunnie, Anna L. E.; Moore, Pippa J.

Published in:
Marine Ecology-Progress Series

Publication date:
2016

The re-use license for this item is:
CC BY-NC-ND

The Document Version you have downloaded here is:
Peer reviewed version

The final published version is available direct from the publisher website at:
[10.3354/meps11544](https://doi.org/10.3354/meps11544)

[Link to author version on UHI Research Database](#)

Citation for published version (APA):
Smale, D. A., Burrows, M. T., Evans, A. J., King, N., Sayer, M. D. J., Yunnie, A. L. E., & Moore, P. J. (2016). Linking environmental variables with regional-scale variability in ecological structure and standing stock of carbon within UK kelp forests. *Marine Ecology-Progress Series*, 542, 79-95. <https://doi.org/10.3354/meps11544>

General rights

Copyright and moral rights for the publications made accessible in the UHI Research Database are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights:

- 1) Users may download and print one copy of any publication from the UHI Research Database for the purpose of private study or research.
- 2) You may not further distribute the material or use it for any profit-making activity or commercial gain
- 3) You may freely distribute the URL identifying the publication in the UHI Research Database

Take down policy

If you believe that this document breaches copyright please contact us at RO@uhi.ac.uk providing details; we will remove access to the work immediately and investigate your claim.

Linking environmental variables with regional-scale variability in ecological structure and standing stock of carbon within kelp forests in the United Kingdom

Dan A. Smale^{1*}, Michael T. Burrows², Ally J. Evans³, Nathan King³, Martin D. J. Sayer^{2,4}, Anna L. E. Yunnice¹, Pippa J. Moore^{3,5}

¹Marine Biological Association of the United Kingdom, The Laboratory, Citadel Hill, Plymouth PL1 2PB, UK

²Scottish Association for Marine Science, Dunbeg, Oban, Argyll, Scotland PA37 1QA

³Institute of Biological, Environmental and Rural Sciences, Aberystwyth University, Aberystwyth, SY23 3DA, UK.

⁴NERC National Facility for Scientific Diving

⁵Centre for Marine Ecosystems Research, School of Natural Sciences, Edith Cowan University, Joondalup 6027, Western Australia, Australia

Table S1. Reported values for carbon content of kelp species, as a percentage of dry weight (list is not exhaustive).

Kelp species	Region	C (% of DW)	No. inds.	Reference
<i>Laminaria digitata</i>	Northern France	29.1	197	(Gevaert et al. 2008)
<i>Laminaria digitata</i>	Nova Scotia	32.7	3	(Mann 1972)
<i>Laminaria digitata</i>	Rhode Island	26.9	3	(Brady-Campbell et al. 1984)
<i>Laminaria hyperborea</i>	Norway	31.3	>32	(Sjøtun et al. 1996)
<i>Laminaria solidungula</i>	Arctic	32.5	>30	(Dunton & Schell 1986)
<i>Saccharina latissima</i>	Rhode Island	31.5	6	(Brady-Campbell et al. 1984)
<i>Saccharina latissima</i>	Northern France	28	182	(Gevaert et al. 2001)
<i>Saccharina longicuris</i>	Nova Scotia	29.3	10	(Mann 1972)
		Mean: 30.16		

Literature cited

- Brady-Campbell MM, Campbell DB, Harlin MM (1984) Productivity of kelp (*Laminaria* spp.) near the southern limit in the Northwestern Atlantic Ocean. 18:79-88
- Dunton KH, Schell DM (1986) Seasonal carbon budget and growth of *Laminaria solidungula* in the Alaskan High Arctic. 31:57-66
- Gevaert F, Davoult D, Creach A, Kling R, Janquin M-A, Seuront L, Lemoine Y (2001) Carbon and nitrogen content of *Laminaria saccharina* in the eastern English Channel: biometrics and seasonal variations. 81:727-734
- Gevaert F, Janquin MA, Davoult D (2008) Biometrics in *Laminaria digitata*: A useful tool to assess biomass, carbon and nitrogen contents. 60:215-219
- Mann KH (1972) Ecological energetics of the seaweed zone in a marine bay on the Atlantic coast of Canada: I. Zonation and biomass of seaweeds. Mar Biol 12:1-10
- Sjøtun K, Fredriksen S, Ruess J (1996) Seasonal growth and carbon and nitrogen content in canopy and first-year plants of *Laminaria hyperborea* (Laminariales, Phaeophyceae). Phycologia 35:1-8

Table S2. Results of multiple linear regressions (DistLM) linking environmental predictor variables with kelp canopy density, canopy biomass and standing stock of carbon. Marginal tests for each predictor variable are shown here, the best most parsimonious models derived from step-wise selection method are presented in Table 4. SS = sum of squares; Prop. = Proportion of variation explained.

(A) Canopy density

Variable	SS	F	P	Prop.
Urchin density	2.1477	0.5980	0.477	0.05
Summer max. temp.	7.971	2.6494	0.14	0.21
Summer daytime light	0.51844	0.1381	0.702	0.01
Depth	0.0078	0.0023	0.982	0.01
Large scale fetch	29.639	35.205	0.001	0.78
Water motion (waves)	21.396	12.842	0.003	0.56
Water motion (tides)	0.96902	0.2612	0.617	0.03
log Chlorophyll a	16.812	7.9136	0.017	0.44
Phosphate	2.4409	0.6853	0.442	0.06
Nitrate+nitrite	2.3998	0.6730	0.416	0.06

(B) Canopy biomass

Variable	SS	F	P	Prop.
Urchin density	0.19703	0.0065	0.951	0.01
Summer max. temp.	114.01	5.9374	0.032	0.37
Summer daytime light	163.08	11.408	0.004	0.53
Depth	12.986	0.4431	0.513	0.04
Large scale fetch	7.3889	0.2474	0.606	0.02
Water motion (waves)	62.927	2.5885	0.116	0.21
Water motion (tides)	71.743	3.0622	0.114	0.23
log Chlorophyll a	30.78	1.1183	0.319	0.10
Phosphate	2.5933	0.0865	0.768	0.01
Nitrate+nitrite	10.312	0.3487	0.567	0.03

(c) Total carbon standing stock

Variable	SS	F	P	Prop.
Urchin density	14539	0.0563	0.843	0.01
Summer max. temp.	818600	4.6055	0.073	0.31
Summer daytime light	1093500	7.2778	0.029	0.42
Depth	52876	0.2079	0.657	0.02
Large scale fetch	99142	0.3970	0.515	0.04
Water motion (waves)	709680	3.7622	0.068	0.27
Water motion (tides)	441010	2.0464	0.181	0.17
log Chlorophyll a	202430	0.8457	0.379	0.02
Phosphate	2504	0.0096	0.915	0.01
Nitrate+nitrite	21441	0.0832	0.775	0.01




Fig S1. Schematic of kelp thallus to show segments obtained for dry versus fresh weight analysis and morphological measurements taken for each canopy-forming sporophyte.




Figure S2. Deployment of light, temperature and water motion sensors in subtidal kelp forests. Data sensors were attached to a sub-surface pellet buoy, which was tethered to the seabed using anchor chain and rope.




Figure S3. Relationships between fresh and dry weight for kelp tissue derived from basal (a) and distal (b) sections of the blade as well as the stipe (c). Biomass data were generated from 47 kelp plants collected from N Scotland, 13 from W Scotland, 48 from SW Wales and 46 from SW England.




Figure S4. Light and temperature conditions recorded during summer 2014 at the study sites. A: Mean daily temperature recorded within each region (average of 3 sites) throughout logger deployment. B: Mean (non-hatched bars) and maximum (hatched bars) daily temperature recorded at each site during 24 overlapping deployment days during summer 2014 (26th July – 18th August). C: Mean daytime light intensity recorded at each study site in summer 2014 (calculated from 14 days of data following logger deployment, based on daytime values only; 0800 to 2000 hours). Site names correspond to those in Figure 1.




Figure S5. Water motion recorded by accelerometers at each study site during summer 2014. Values are estimated ms^{-1} recorded during 45 day deployment of sensor array. Tidal cycles derived from sea level readings from each study region are also shown. Data were obtained from the following tidal gauges: N Scotland (sites A1-A3): Wick; W Scotland (sites B1-B3): Tobermory; SW Wales (sites C1-C3): Milford Haven; SW England (sites D1-D3): Devonport (sea level data are managed and distributed by the British Oceanographic Data Centre, www.bodc.ac.uk). Site names correspond to those in Figure 1.




Fig. S6. Relationship between the metric derived to quantify tidal motion (from *in situ* accelerometers) and range in sea level for each 12-hour period during the 45-day sensor deployment ($n = 90$). Data are shown for 2 representative sites, North Grahamsay in northern Scotland (site A3) and Stoke Point in southwest England (site D2). As expected, there was a positive relationship between the variables, with periods of high diurnal sea level variability (i.e. during spring tides) corresponding with high values of water motion induced by tidal flow.