

## UHI Research Database pdf download summary

### **A rare occurrence of reversal in the common megrim *Lepidorhombus whiffiagonis* (Pleuronectiformes: Scophthalmidae) in the northern North Sea**

Macdonald, Paul

*Published in:*  
Journal of Fish Biology

*Publication date:*  
2013

*The re-use license for this item is:*  
CC BY-NC

*The Document Version you have downloaded here is:*  
Early version, also known as pre-print

*The final published version is available direct from the publisher website at:*  
[10.1111/jfb.12204](https://doi.org/10.1111/jfb.12204)

### [Link to author version on UHI Research Database](#)

*Citation for published version (APA):*

Macdonald, P. (2013). A rare occurrence of reversal in the common megrim *Lepidorhombus whiffiagonis* (Pleuronectiformes: Scophthalmidae) in the northern North Sea. *Journal of Fish Biology*, 83(3), 691-694. <https://doi.org/10.1111/jfb.12204>

#### **General rights**

Copyright and moral rights for the publications made accessible in the UHI Research Database are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights:

- 1) Users may download and print one copy of any publication from the UHI Research Database for the purpose of private study or research.
- 2) You may not further distribute the material or use it for any profit-making activity or commercial gain
- 3) You may freely distribute the URL identifying the publication in the UHI Research Database

#### **Take down policy**

If you believe that this document breaches copyright please contact us at [RO@uhi.ac.uk](mailto:RO@uhi.ac.uk) providing details; we will remove access to the work immediately and investigate your claim.

1 **A rare occurrence of reversal in the common megrim, *Lepidorhombus***  
2 ***whiffiagonis* (Walbaum, 1792) (Pleuronectiformes: Scophthalmidae), in the**  
3 **northern North Sea**

4

5

6 Macdonald, P.

7

8

9 NAFC Marine Centre, Marine Science & Technology Department, Port Arthur,  
10 Scalloway, Shetland, ZE1 0UN, Scotland, UK.<sup>1</sup>

11

12

13 Rare occurrence of reversal in common megrim

14

15

16

17

18

19

20

21

22

23

24

---

<sup>1</sup> Tel.: +44 01595 772331; fax: 01595 772001; email: paul.macdonald@nafc.uhi.ac.uk

25 **Abstract**

26 An adult (231 mm  $L_S$ ; 371 mm  $L_T$ ), dextral (right-sided) common megrim,  
27 *Lepidorhombus whiffiagonis*, is reported from the northern North Sea. This is the first  
28 recorded occurrence of reversal in this species. Other than its reversed asymmetry,  
29 the specimen had similar morphological and meristic characteristics to those of non-  
30 reversed individuals of the same species.

31

32

33 Keywords: flatfish, sinistral, dextral, reversed asymmetry

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50 Ontogenetic metamorphosis in flatfishes (Pleuronectiformes) results in one eye  
51 migrating from one side of the head to the other (Brewster, 1987). The side to which  
52 the eye migrates, resulting in a left-sided (sinistral) or right-sided (dextral) orientation,  
53 is typically species dependent. However, in the most primitive flatfishes, the  
54 Psettodidae (spiny turbot), the eye migrates to the left or right side with equal  
55 frequency (Janvier, 2008). For many species of flatfishes, the majority of individuals  
56 exhibit the same orientation, although occurrences of reversal, resulting in sinistral  
57 orientation in dextral species and *vice versa* have been well documented (Norman,  
58 1934; Ahlstrom *et al.*, 1984). Occurrences of reversal have been reported in a broad  
59 range of species of flatfishes and in some species it is commonly encountered  
60 (Norman, 1934; Gartner, 1986). Studies on the left-sided Californian flounder  
61 *Paralichthys californicus* (Ayres 1859) have reported that the incidence of reversal in  
62 this species is as high as 40% (Ginsburg, 1952; Kramer *et al.*, 1995). Hubbs and  
63 Kuronuma (1942) noted that starry flounder *Platichthys stellatus* (Pallas 1787) exhibit  
64 a geographical cline in asymmetry with sinistrality increasing from 50% in California to  
65 100% in Japan. However, cases of reversal are typically rarer for the majority of  
66 species of flatfishes. For example, Bruno and Fraser (1987) reported a single case of  
67 reversal amongst 15,859 common dab *Limanda limanda* (L. 1758) sampled in the  
68 North Sea. Similarly, occasional occurrences of reversal have been reported in  
69 summer flounder *Paralichthys dentatus* (L. 1766) (Gudger, 1936), Atlantic halibut  
70 *Hippoglossus hippoglossus* (L. 1758) and plaice *Pleuronectes platessa* (L. 1758)  
71 (Gudger, 1935).

72

73

74 The common megrim *Lepidorhombus whiffiagonis* (Walbaum 1792) (Scophthalmidae)  
75 is a commercially important flatfish distributed from Iceland to the Mediterranean Sea  
76 (Nielsen, 1989). It is one of two species of the genus *Lepidorhombus* (Günther 1862),  
77 the other being the four-spotted megrim *Lepidorhombus boscii* (Risso 1810). One  
78 adult (231 mm  $L_S$ ; 371 mm  $L_T$ ) reversed common megrim was caught 100 km  
79 northwest of the Shetland Isles in June 2009 during sampling of megrim in the northern  
80 North Sea (Figure 1). The specimen is a mature male and was captured using a  
81 demersal twin-rig otter trawl at a depth of 196 m. It represents the only case of reversal  
82 recorded for 39,072 measured individuals of this species that were captured in the  
83 North Sea between May 2008 and March 2012. In the reversed individual, interorbital  
84 width is less pronounced and caudal peduncle length is greater (Table I) than  
85 corresponding values of non-reversed individuals. No other significant variation in  
86 meristic features or patterning and coloration were apparent in this individual,  
87 suggesting that reversal did not result in significant changes in the external  
88 morphology and colouration of this individual. In contrast, a similar study of reversal in  
89 the shallow-water flounder *Paralichthys orbignyanus* (Valenciennes 1839) reported  
90 several differences in a number of comparative measurements, including head,  
91 maxilla and prepectoral lengths, between non-reversed individuals and a reversed  
92 specimen of this species (Diaz de Astarloa, 1997).

93

94

95 Reversals in megrims are not limited to *L. whiffiagonis*. Vassilopoulou (1993) reported  
96 a small number of occurrences (0.026%) of reversed (dextral) individuals from  
97 samples of *L. boscii* collected in the Aegean Sea, indicating that reversal exists, but is  
98 rare in species of *Lepidorhombus*.

99

100

101 This is the first reported occurrence of reversal in common megrim. Due to the rarity  
102 of reversed specimens in this species, it has been preserved and deposited in the fish  
103 collection of the NAFC Marine Centre, Scalloway, Shetland. The collection is not  
104 formally catalogued and therefore does not assign numbers to specimens.

105

106

107 Acknowledgements

108 This study was carried out during a wider investigation into the biology and ecology of  
109 megrim in the northern North Sea and was partly funded by the Seafish Industry  
110 Authority, Scottish Fishermen's Trust and Shetland Islands Council. I am grateful to  
111 Marine Scotland-Science and all the fishermen and who provided access to their  
112 vessels for sampling. I am also grateful to two anonymous reviewers for their insightful  
113 comments.

114

115

116 **References**

117 Ahlstrom, E. H., Amaoka, K., Hensley, D. A., Moser, H. G. & Sumida, B. Y. (1984).  
118 Pleuronectiformes: development. In *Ontogeny and Systematics of Fishes* (Moser, H.  
119 G., Richards, W. J., Cohen, D. M., Fahay, M. P., Kendall, W., Jr. & Richardson, S. L.,  
120 eds.), pp. 640-670: Special Publication 1. American Society of Ichthyologists and  
121 Herpetologists.

122 Brewster, B. (1987). Eye migration and cranial development during flatfish  
123 metamorphosis: a reappraisal (Teleostei: Pleuronectiformes). *Journal of Fish Biology*  
124 **31**, 805-833.

125 Bruno, D. W. & Fraser, C. O. (1987). A case of reversal in the common dab, *Limanda*  
126 *limanda* (L.). *Journal of Fish Biology* **32**, 483-484.

127 Diaz de Astarloa, J. M. (1997). A case of reversal in *Paralichthys orbignyanus* a  
128 shallow-water flounder from the south-western Atlantic. *Journal of Fish Biology* **50**,  
129 900-902.

130 Gartner, J. V. (1986). Observations on anomalous conditions in some flatfishes  
131 (Pisces: Pleuronectiformes), with a new record of partial albinism. *Environmental*  
132 *Biology of Fishes* **17**, 141-152.

133 Ginsburg, I. (1952). Flounders of the genus *Paralichthys* and related genera in  
134 American waters. *Fishery Bulletin* **52**, 267-351.

135 Gudger, E. W. (1935). Abnormalities in flatfishes (Heterosomata). I. Reversal of sides:  
136 a comparative study of the known data. *Journal of Morphology* **58**, 1-39.

137 Gudger, E. W. (1936). A reversed almost wholly ambicolorate summer flounder,  
138 *Paralichthys dentatus*. *American Museum Novitates* **896**, 1-5.

139 Hubbs, C. L. & Kuronuma, K. (1942). Hybridization in nature between two genera of  
140 flounders in Japan. *Papers of the Michigan Academy of Science, Arts and Letters* **27**,  
141 267-306.

142 Janvier, P. (2008). Palaeontology: Squint of the fossil flatfish. *Nature* **454**, 169-170.

143 Kramer, D. E., Barss, W. H., Paust, B. C. & Bracken, B. E. (1995). Guide to Northeast  
144 Pacific flatfishes: Families Bothidae, Cynoglossidae, and Pleuronectidae. In *Marine*  
145 *Advisory Bulletin # 47*, p. 104: Alaska Sea Grant College Program, University of  
146 Alaska, Fairbanks, Alaska.

147 Nielsen, J. G. (1989). Scophthalmidae. In *Fishes of the North-eastern Atlantic and the*  
148 *Mediterranean (FNAM)* (Whitehead, P. J. P., Bauchot, M.-L., Hureau, J.-C., Nielsen,  
149 J. & Tortonese, E., eds.), pp. 1286-1293. Unesco, Paris: Springer-Verlag, Berlin.  
150 Norman, J. R. (1934). A systematic monograph of the Flatfishes (Heterosomata). Vol  
151 1: Psettodidae, Bothidae, Pleuronectidae. 459 p. London: British Museum (Nat. Hist.).  
152 Vassilopoulou, V. (1993). Orientation anomalies (reversal) of the four-spotted megrim  
153 (*Lepidorhombus boscii*) in the Aegean Sea. *Journal of Fish Biology* **45**, 165-166.

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168



169 Table 1 A comparison of morphological features of a reversed and non-reversed  
 170 *Lepidorhombus whiffiagonis*.

	Reversed	Non-reversed <i>L. whiffiagonis</i> (n=22)		
	<i>L. whiffiagonis</i>	Range	Mean	S.D.
Total length ( $L_T$ )	331	314 - 357	329	10.8
Standard length ( $L_S$ )	272	262 - 302	276	11.6
Head length	28.3	27.2 - 30.3	28.5	3.59
Prepectoral distance	27.2	26.8 - 29.6	27.9	3.34
Maxilla length	16.5	15.2 - 17.5	16.3	1.78
Caudal peduncle length	20.6	14.3 - 20.1	18.4	3.45
Pectoral fin length	16.1	11.6 - 18.4	15.1	3.05
Interorbital width	0.3	0.5 - 1.0	0.7	0.29
Dorsal-fin rays	90	84 - 93	87	2.28
Anal-fin rays	67	65 - 72	68	2.27
Pectoral-fin rays	11	10 - 12	11	0.45

171 Total and standard lengths are reported in mm: all other measurements are % standard length

172

173


174

175

176

177

178



179

180 Figure 1 Reversed specimen of *Lepidorhombus whiffiagonis*, 272 mm Ls (top), and  
181 non-reversed specimen, 284 mm Ls (bottom), caught in the northern North Sea.

182

183

184

185

186