

UHI Research Database pdf download summary

Turning off the DRIP ('Data-rich, information-poor') - rationalising monitoring with a focus on marine renewable energy developments and the benthos

Wilding, Tom; Gill, Andrew B.; Boon, Arjen; Sheehan, Emma; Dauvin, Jean-Philippe; Pezy, Jean-Philippe; O'Beirn, Francis; Janas, Urszula; Rostin, Liis; De Mesel, Ilse

Published in:
Renewable and Sustainable Energy Reviews

Publication date:
2017

Publisher rights:
© 2017 SAMS. Published by Elsevier Ltd.

The re-use license for this item is:
CC BY-NC-ND

The Document Version you have downloaded here is:
Publisher's PDF, also known as Version of record

The final published version is available direct from the publisher website at:
[10.1016/j.rser.2017.03.013](https://doi.org/10.1016/j.rser.2017.03.013)

[Link to author version on UHI Research Database](#)

Citation for published version (APA):

Wilding, T., Gill, A. B., Boon, A., Sheehan, E., Dauvin, J., Pezy, J., O'Beirn, F., Janas, U., Rostin, L., & De Mesel, I. (2017). Turning off the DRIP ('Data-rich, information-poor') - rationalising monitoring with a focus on marine renewable energy developments and the benthos. *Renewable and Sustainable Energy Reviews*, 74, 848-859. <https://doi.org/10.1016/j.rser.2017.03.013>


General rights

Copyright and moral rights for the publications made accessible in the UHI Research Database are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights:

- 1) Users may download and print one copy of any publication from the UHI Research Database for the purpose of private study or research.
- 2) You may not further distribute the material or use it for any profit-making activity or commercial gain
- 3) You may freely distribute the URL identifying the publication in the UHI Research Database

Take down policy

If you believe that this document breaches copyright please contact us at RO@uhi.ac.uk providing details; we will remove access to the work immediately and investigate your claim.



Turning off the DRIP ('Data-rich, information-poor') – rationalising monitoring with a focus on marine renewable energy developments and the benthos



Thomas A. Wilding^{a,*}, Andrew B. Gill^b, Arjen Boon^c, Emma Sheehan^d, Jean-Claude Dauvin^e, Jean-Philippe Pezy^e, Francis O'Beirn^f, Urszula Janas^g, Liis Rostin^h, Ilse De Meselⁱ

^a SAMS, Scottish Marine Institute, OBAN, Scotland PA37 1QA, UK

^b Centre for Offshore Renewable Energy and Engineering, School of Water, Energy, and, Environment, Building 52, Cranfield University, Cranfield, Bedfordshire MK43 0AL, UK

^c Deltares, Delft, The Netherlands

^d Plymouth University Marine Institute, School of Biological and Marine Sciences, Drake Circus, Plymouth PL4 8AA, UK

^e Normandie Université, UNICAEN, UMR M2C (UCN, UR, CNRS-6143), 24 rue des Tilleuls, 14000 Caen cedex 5, France

^f Marine Institute, Rinville, Oranmore, Galway, Ireland

^g Institute of Oceanography, University of Gdańsk, Al. Marsz. Piłsudskiego 46, Gdynia, Poland

^h Estonian Marine Institute, University of Tartu, Mäealuse 14, 12618 Tallinn, Estonia

ⁱ Royal Belgian Institute of Natural Sciences (RBINS), Operational Directorate Natural Environment, Marine Ecology and Management, 3e en 23e Linierregimentsplein, 8400 Oostende, Belgium

ARTICLE INFO

Keywords:

Benthos
Biodiversity
Confidence and risk
Environmental impact assessment
Monitoring
Spatial and temporal scales
Study design
Offshore
Tidal- and wave-power
Wind-turbines

ABSTRACT

Marine renewable energy developments (MREDs) are rapidly expanding in size and number as society strives to maintain electricity generation whilst simultaneously reducing climate-change linked CO₂ emissions. MREDs are part of an ongoing large-scale modification of coastal waters that also includes activities such as commercial fishing, shipping, aggregate extraction, aquaculture, dredging, spoil-dumping and oil and gas exploitation. It is increasingly accepted that developments, of any kind, should only proceed if they are ecologically sustainable and will not reduce current or future delivery of ecosystem services. The benthos underpins crucial marine ecosystem services yet, in relation to MREDs, is currently poorly monitored: current monitoring programmes are extensive and costly yet provide little useful data in relation to ecosystem-scale-related changes, a situation called 'data-rich, information-poor' (DRIP). MRED –benthic interactions may cause changes that are of a sufficient scale to change ecosystem services provision, particularly in terms of fisheries and biodiversity and, via trophic linkages, change the distribution of fish, birds and mammals. The production of DRIPy data should be eliminated and the resources used instead to address relevant questions that are logically bounded in time and space. Efforts should target identifying metrics of change that can be linked to ecosystem function or service provision, particularly where those metrics show strongly non-linear effects in relation to the stressor. Future monitoring should also be designed to contribute towards predictive ecosystem models and be sufficiently robust and understandable to facilitate transparent, auditable and timely decision-making.

1. Introduction

Harnessing marine renewable energy resources (e.g. wind, wave, tidal stream) at commercial scales, involves deployment of multiple devices and associated infrastructure within a defined area of the coastal or offshore environment. Worldwide the Marine Renewable Energy (MRE) sector is growing significantly in line with technological development and the political imperative to act on climate change. The

predicted 3.6 GW of capacity to be installed within Europe by 2030 and potentially 188 GW (~50,000 wind-turbines) by 2050 illustrates the scale of development (European Ocean Energy Association 2010).

MRE Developments (MREDs) are part of existing and ongoing modification of our seas and seabed that are occurring at local to regional scales that include commercial fishing, transport, aggregate extraction, aquaculture, dredging, spoil-dumping and hydrocarbon exploitation. The goods and ecosystem services already derived from

* Correspondence author.

E-mail address: tom.wilding@sams.ac.uk (T.A. Wilding).

coastal waters and seas are substantial [1] and the expansion of MREDS will cause anticipated and unanticipated changes [2–4], that will, to some extent, affect ecosystem service value and capacity. Whilst MREDS will contribute to a reduction in greenhouse gas emissions, there is a clear need to ensure future effective management of the marine environment. Effective MRED management will occur in the light of potentially cumulative impacts and necessitates appropriately-scaled and focussed monitoring to ensure that the inevitable changes attributable to them are acceptable to society.

Monitoring marine developments, to assess their impacts on the receiving environment, is a requirement under many regulations and regulatory frameworks (e.g. numerous European Directives, US FDA legislation, Australian EPBC Act, 1999). Within Europe, these regulations are being rigorously tested by the extensive plans for MREDS. Monitoring of the environmental effects of MREDS is currently split between four broad receptor groups - mammals, birds, fish¹ and benthos. Attention has tended to focus on iconic/charismatic species groups (marine mammals and birds) mainly because such groups have both high public acclaim [5] and protection (e.g. under the EU Birds and Habitats Directives [3]) and there are demonstrable examples of them being affected directly by MREDS [6].

In accordance with ecosystem-based thinking and recent environmental legislation, monitoring requirements are shifting from a species-centric focus towards understanding and managing the marine environment in an integrative, holistic manner (e.g. the Marine Strategy Framework Directive (MSFD) in Europe) with due consideration given to the cumulative impacts of activities across various sectors [7]. Consequently, environmental impacts need to be assessed at ecosystem- rather than local-scales [8] which requires a profound understanding of the benthic components of the ecosystem owing to its pivotal role in delivering and supporting key ecosystem services [9]. Any MRED-related changes to the benthic ecosystem, even if hard to detect, may still have profound implications for the provision of valuable ecosystem services, including those related to mammals, birds and fish, because of non-linear effects [10,11].

Ineffective quantification of ecosystem-function change associated with the MREDS-benthos interaction compromises our ability to determine whether any predicted/measured change is deemed meaningful,² where meaningful is defined as changes that affect the ecological components fundamental to ecosystem service provision [12] at any logically justified scale. In order to assess meaningful change, within a limited budget, it is necessary to define acceptable limits of change (targets/thresholds) associated with populations and/or functional attributes (e.g. biomass production or biogeochemical processing). In any monitoring assessment, consideration should be given to the probability that the thresholds have been exceeded and, if necessary, recommend appropriate management actions or mitigation measures.

The magnitude and extent of human-impacts that are relevant to quantifying MRED-induced ecological change requires an in-depth understanding of ecological process occurring at a range of scales [13]. Despite the considerable expenditure in MRED-benthic monitoring programmes the impact of MREDS on ecosystems, as propagated via the benthic ecosystem, remains poorly understood [3]. In relation to MREDS many existing benthos-relevant research /mapping programmes have lacked clarity and rigour [14], have been formulaic ('box-ticking'), unrelated to justified temporal or spatial scales [6] and, as a result, have not enhanced an understanding of MRED interactions

¹ In this context, 'fish' refers to all fish, and shellfish, with a focus on those of commercial and conservation importance including migratory /diadromous species such as eels and salmonids.

² The term 'meaningful', used here to mean a change that any stakeholder cares about, is deliberately subjective. The alternative 'significance' is not used because it should always be clarified as to whether the reference is to 'statistical', 'ecological' or other significance.

at relevant ecosystem scales [15]. This situation is known by the acronym DRIP- 'data-rich, information-poor' [16].³ As a consequence, current MRED-associated benthic monitoring programmes have difficulty in raising legitimate concerns or confirming that unacceptable levels of change are highly unlikely to be occurring. Such costly monitoring programmes thus add to the 'DRIP' [15,17] and also contribute to delays in the mandatory Environmental Impact Assessment (EIA) process which adds further cost burden on the whole sector [18]. MREDS include a number of technologies (e.g. wind, wave and tide) of which only offshore wind has been widely commercialised⁴ and which, consequently, forms the focus of this review. Whilst offshore tide and wave developments will be located in very different receiving environments compared with offshore wind the core issues raised in this review are directly relevant to all sectors.

The objectives of this paper are to:

- Highlight the need for clear management objectives in deriving a logical monitoring programme
- Review benthic metric selection and the need to set thresholds around which change can be logically assessed
- Consider relevant scales over which to assess change
- Review how and why 'DRIPy' data are currently generated and promote an alternative to null hypothesis significance testing as the basis for monitoring
- Propose a 'traffic-light' method for decision making in relation to thresholds
- Review current practices in assessing change in benthic environments
- Propose a structured approach to data-gathering to curtail the generation of DRIPy data
- Promote the re-direction of currently wasted resources to generating data which is useful to assessing relevant ecosystem-level change

Through our analyses, we look to promote a step change in the basis for environmental monitoring of MREDS that is applicable to any receptor group but we focus on the benthos because of its key and under-considered ecosystem role [19].

2. Monitoring MRED-benthic interactions

The need to include the benthos within ecosystem-led monitoring is fundamental [20,21]. The current problems associated with existing MRED-benthic monitoring programmes stem from a number of interlinked issues including the lack of clear management objectives (undefined scales, thresholds and metrics), fundamental problems with data gathering and interpretation and the technical/logistical problems associated with data collection.

2.1. Management objectives

Monitoring programmes, of any kind, can only be properly designed and successful where the objectives are clearly specified [22] and, in respect of MREDS, these are frequently lacking [14]. Without clear objectives, monitoring becomes a cataloguing exercise and, whilst this may have use in terms of ad-hoc future research programmes, such cataloguing is not cost-effective in development-specific studies on which to base management decisions. Objectives need to be clear and targeted [23].

In most nations, maritime activities are controlled via a number of regulatory measures that are specified at a range of political levels [24]. Within the EU the main drivers of monitoring in relation to the

³ In this context 'DRIPy data' are characterised by being 'data-rich yet information poor'.

⁴ See Maygen (<http://www.meygen.com/the-project/>) for an example of a commercial-scale tidal array in construction.

ecological effects of MREDS centre around the Birds Directive (2009/147/EC), the Habitats Directive (92/43/EEC) and the Environmental Impact Assessment Directive (including cumulative impact assessment) [7,17,25]. These directives are variously transposed into Member States' (MS) national legislative framework and, consequently, there is limited common (EU) basis for monitoring around MREDS. Within Europe, the monitoring-emphasis is on iconic taxa such as marine mammals and birds or specially protected habitats (e.g. reef-forming species such as *Sabellaria spinulosa* (Leuckart, 1849)). Whilst Directives such as the Water Framework Directive [12] and Marine Strategy Framework Directive outline ecosystem-level criteria (e.g. 'good-ecological status'⁵ and 'good environmental status'⁶ respectively [26,27] these are not applied in relation to specific developments. However, the WFD can be applied to specific marine features, of indeterminate scale, which could encompass a specific renewable development (see §2.3). Whilst cumulative impact assessments are required for MREDS (e.g. as part of EIA in the EU) these are often lacking [28] largely because the underlying principles and definitions are poorly specified or understood [7,25,29]. Within the EU, the ecosystem-service-maintaining goals of Directives such as the MSFD have to be balanced against the 'Blue Growth' agenda (e.g. under the Integrated Maritime Plan) [24] and within the context of marine spatial planning (under the Maritime Spatial Planning Directive) [30]. Across the EU there is a lack of strategic direction, both within and between Member States, in terms of understanding benthic-ecosystem-industry interactions [31] and this contrasts, at least in part, with other receptor groups.

Of the four commonly assessed receptor groups (mammals, birds, fish and benthos) the benthic-MRED interaction has been the least studied. For marine mammals and birds, management objectives are better articulated when compared with the benthos and the most logically justified monitoring objectives focus on the long-term sustainability of populations. Whilst there remain numerous data-gaps [32], many marine mammal and bird populations are *relatively* well understood in terms of their basic life-traits such as fecundity, age-to-maturity, natural mortality rates, spatial distribution and population mixing and boundaries [32,33]. For many species there exists extensive time-series data [34] and recently these have been augmented using data from tracking devices attached to individual animals [35]. Population sustainability in the face of developments, such as MREDS, has been assessed using models (e.g. Potential Biological Removal/Potential Consequences of Disturbance models and/or Population Viability Analyses) that are beginning to capture some of the complexity inherent in natural populations [6,14,32,33,35–37]. The issue of uncertainty, in assessing extinction risk, has also been addressed in relation to some marine mammals and birds, identified under the auspices of the International Union for Conservation of Nature [38].

Fisheries management objectives typically include maintaining stocks near their maximum sustainable yield and, again, many fisheries species are *relatively* well understood in terms of their fecundity and natural mortality [39] and likely non-linear relationships between stocks and some environmental drivers [40]. Historical fisheries data

are available and give an indication of long-term trends in various populations (as a function of fishing intensity and environmental parameters) and modern genomics techniques have indicated the location and nature of some stock boundaries [39]. In addition, within Europe, many fish-stocks are statutorily monitored and afforded protection under the Common Fisheries Policy [24] which explicitly acknowledges the importance of defining spatial boundaries in respect of fishery management. In contrast, knowledge of benthic-ecosystem or benthic-population-scale processes is often lacking and/or not used in informing and justifying rational approaches to assessing MRED-benthic interactions. This is at odds with the ecosystem-approach to assessing change in relation to man's activities in the oceans [8] and, within Europe, Directives such as the MSFD and the cumulative impact requirement under the EIA Directive.

MRED-benthos monitoring programmes around existing developments have tended to focus on three aspects: (i) macrobenthic species richness, abundance/biomass (via grab-sampling in sedimentary environments [14,17,41,42]), (ii) epibenthic cover/diversity (on hard substrata; e.g. [42–45] or (iii) epibenthic diversity as determined from trawls [42]. These three aspects are usually assessed at arbitrary spatial scales, are descriptive [42] and not linked to ecosystem-service provision. We believe that the most relevant benthic-related ecosystem services to be affected by changes in the benthos around MREDS will be related to fisheries [46], trophic networks (potentially affecting top-predators such as fish, birds and marine mammals) [32,35], sediment-based nutrient processing (biogeochemical reactor) and biodiversity [4].

2.2. Benthic metric selection and thresholds

The choice of metric (response variable) in benthic monitoring programmes is a major consideration and will determine the methodology employed, the appropriate spatial- and temporal-domain (§2.3) and the necessary confidence required in any assessment (§2.4.2). In general terms, whatever metric is chosen needs to be easily/cheaply measurable, sensitive to change and specific to the source of impact, predictable and anticipatory (able to signify impending change) [47]. In addition, the chosen metric should be relevant to assessing change that can be averted by management-actions, integrative (i.e. can indicate change over key gradients), have low variability (§2.5), be transferable (to various systems) and be clear and understandable [47]. We would add that metrics should be societally/stakeholder- and ecosystem-service related. Hattam et al. [47] identified a wide-range of ecosystem services delivered by the Dogger Bank (North Sea), and proposed a number of metrics for the assessment of these services. These metrics including food provision (annual carbon production and monetary value), waste-treatment and assimilation (as measured by a healthy benthos in terms of biodiversity, §2.3.3), migratory and nursery habitat provision (as measured by the area of suitable habitat, quality of habitat, number of target species and occupancy rates) among many others.

Monitoring programmes that are based around a question such as 'is there any change in X as a consequence of the development' are limited (§2.4); monitoring should occur in relation to thresholds with a threshold being defined as a "target level or state based on the avoidance of unacceptable outcomes or an ecologically defined shift in system status" [48]. Identifying suitable metrics and negotiating threshold values, which meet the above criteria is difficult, potentially requiring multi-stakeholder consultation [49], and may constitute a significant part (in time and resources) of any monitoring programme [50,51]. The requirement for quantitative regulatory thresholds in respect of the benthos is, however, already part of the EU MSFD and WFD and EU member states have developed numerous benthic-diversity-based metrics [52–54] to classify the ecological status of sediments. The thresholds developed under the WFD offer some potential, at least conceptually, to MRED monitoring, but have been

⁵ The Water Framework Directive (WFD) classifies ecological status on the basis of a number of ecological (including benthic macrofauna) and physio-chemical characteristics. The Directive is comprehensive in terms of the composite characters for each of the indicators used: benthic status, for example, is assessed by comparing diversity, abundance and sensitivity metrics against baseline conditions (see main text). Metrics used in the WFD score between 0 ('Bad') and 1 ('High') with intermediate values of 'Poor', 'Moderate' and 'Good' [52]: there are clear thresholds.

⁶ The MSFD specifies eleven descriptors, covering a suite of pressures and environmental state receptors. The MSFD descriptor 6 (Seafloor Integrity) can be applied to the benthos; it currently contains two indicator groups and covers the physical state of the habitat (biogenic reef extension, and extent human physical disturbance), and benthic biological diversity. There remains considerable variability in the definition of 'good' between MS [12]

beset by problems in defining appropriate baselines.⁷ Identifying and maintaining ecosystem-service-provision maybe a more tractable objective.

In the European Union, under the MSFD, there is a requirement to achieve ‘good environmental status’ (GES) for all parts of the ecosystem and there are clear objectives in relation to benthos (but see [27]). GES relates to ecosystem health which is defined as ‘the condition of a system that is self-maintaining, vigorous, resilient to externally imposed pressures, and able to sustain services to humans ...’ [13]. Quantifying ecosystem health is problematic, particularly when faced with an unknown baseline, but one possible mechanism is to examine time-series based ‘ecosystem trajectories’ [13], possibly in relation to changes in the ratio between macrobenthic components [55] or changes in average regional diversity [20]. Ecosystem models, coupled with earth-observations may indicate potential candidate benthic species/groups and thresholds for monitoring MREDS at larger (e.g. regional sea) scales [56].

In relation to all monitoring programmes, threshold setting will be easiest (most justifiable) where the monitored metric is known or predicted to respond in a strongly non-linear way to the development (e.g. in fisheries [40]). Predicting non-linear effects in the face of numerous and complex ecological interactions that will occur following large-scale MRED deployment is difficult [26,57,58]. However, if identified, a strongly non-linear response may indicate the presence of a ‘tipping-point’ around which management decisions (with safety margins) are relatively easily argued and justified to stakeholders [11,26]. In Scotland (UK), regulators have specified limits on species reductions and various chemical parameters occurring around point-sources of impact (e.g. fish-farms [59]). These limits are set around tipping points and ensure that the benthic system’s assimilative processes can continue [59] and are thus logically based. Benthic-related thresholds for heavy metals are specified, in relation to aquaculture, and also form part of Environmental Statements in offshore renewables developments e.g. [60] and have been applied in other contaminated sites e.g. Puget Sound, Washington, US [61] but the rationale for the spatial scale of any assessment is not necessarily clear (§2.3).

Whilst wind-turbines and their supporting infrastructure (including scour protection material) may occupy > 1000 m² per device [62] it is unknown whether this degree of modification (multiplied by array size) would cause meaningful changes to ecosystem-functioning or service provision. Whilst impacts of a single device will be irrelevant on the ecosystem-scale, studies based on individual structures can be related to ecosystem-service provision by aiding understanding of trophic linkages. For example, the data on size-specific cod growth rates and fidelity (which constitute the metrics under investigation) around individual offshore wind-turbine foundations [46] is relevant where it is used to inform models that predict regional-scale change in fish-population or behaviour. Understanding such population-scale change helps explain the attractiveness of MREDS to foraging mammals [35] and birds [63] and informs regulators in applying more species-specific regulations (e.g. under the Birds and Habitats Directive in the EU). Whatever metrics and thresholds are chosen they must be assessed at scales that are relevant to the questions being asked.

⁷ In any ecological assessment, change has to be compared with a baseline. When monitoring is initiated the observed baseline condition is often misunderstood to be the ‘natural’ state rather than the ‘current’ state (‘Shifting Baseline Syndrome’ [131]). Shifting, or unknown, baselines are particularly pertinent to coastal zones which have frequently been exposed to intense disturbance [132,133] and issues around baseline-identification have caused problems in the implementation of the WFD [53,134]. Care should be taken to ensure that development objectives, of showing no ‘significant’ local-scale change, over baseline conditions, meet the longer-term, larger-scale objectives as part of ecosystem-scale assessment.

2.3. Relevant scales in relation to MRED-benthic interactions

Designating appropriate scales for monitoring programmes is a challenge particularly when assessing change at lower trophic levels where population boundaries are often poorly defined/understood or highly variable and where species are migratory and/or exhibit considerable ontogenetic changes in habitat utilisation (e.g. have a planktonic stage). What constitutes an appropriate scale to which inference should be made will, in part, be determined by the relevant regulatory framework: these may include locally agreed scales e.g. bays, inlets or specific features⁸ to regional, sea-scale multinational agreements (e.g. EU MSFD). From an ecosystem perspective geopolitical boundaries may be useful (e.g. MSFD Baltic and Celtic Sea sub-regions) but only where they coincide with eco/hydrologically defined boundaries that are relevant to the distribution of the species under investigation. In the UK, sea-areas have been defined by temperature, depth and current [64] and these with, potentially, any of the delineations as defined under the Marine Spatial Planning Initiative (a global initiative with ~20 participating nations [65]) may also be logical in terms of defining spatial limits to benthic monitoring.

2.3.1. Structure colonisation and scale of change

Many MREDS offer novel, surface piercing, hard and, at some scales, structurally complex, surfaces that may make a substantial relative contribution to this type of habitat over national or regional scales [66]. Where the new surface enables the local growth of large populations, particularly of ecosystem engineers (e.g. suspension feeders such as mussels [28]) or non-indigenous species then ecosystem-scale changes should be assessed. Mussels (in culture) have been shown to deplete overall plankton biomass [67], change planktonic communities [68,69] and enhance the reproductive success of their predators with potentially large-scale consequences [70]. Joschko et al. [71] predicted that a biomass of 265 kg (ash-free dry weight) of the blue mussel (*Mytilus edulis* L., 1758) would develop per wind turbine in the German Bight. When extrapolated across an 80-turbine array in the area, it was predicted that there would be a 10% increase in mussel biomass that would consume 1.4% of the annual primary production in that region [71]. Whether this, or similar cumulative effects from future planned developments, would constitute an ‘ecosystem-relevant’ change, within this spatial domain, depends on perspective and requires further research. MRED structures have also been associated with changes in jellyfish populations in the Baltic [72] by providing a hard-substratum for the asexual benthic phase (an effect also seen in respect of aquaculture structures in Thailand [73]) whilst artificial reefs have facilitated the spread of a harmful dinoflagellate, via its benthic stage, in large parts of the Gulf of Mexico [74]. Relevant scales for monitoring in these cases would be the individual structures (e.g. devices) but the results would need to be relevant within a region-scale context necessitating a profound understanding of the ecology of the species concerned and the receiving environment [56,58,75]. At certain scales changes in plankton (e.g. increased jelly-fish populations) may be a nuisance in relation to amenity use and/or damaging to other stakeholders (e.g. the aquaculture sector [76,77]) so any monitoring programme should link potential effects to other activities occurring within relevant spatial domains.

2.3.2. Non-indigenous species

MREDS may facilitate movement of non-indigenous and indigenous species into new habitats [2]. Movement of organisms between devices will occur in a number of ways including, for those with a benthopelagic life-history, the use of MRED devices as ‘stepping stones’ as

⁸ For example, St. Catherine’s Deep, an unusual channel of ~10 x 2 km located on the south side of the Isle of Wight, UK and a potential tidal-development site. This locally unique feature might justifiably form its own spatial boundary within which monitoring should be focused.

seen in relation to oil and gas platforms [78] and which is dependent on the connectivity between the structures [45,79]. Other species may use construction and maintenance vessels on which to ‘hitch-a-ride’ [2]. This means that different deployment patterns /spatial distributions of devices, over regional scales, will have different impacts even whilst the overall number of devices remains constant. Location can play an important part in determining the prevalence of non-indigenous species (NIS), for example, the intertidal foundations of offshore windfarms have been associated with a much greater incidence of NIS compared with sub-tidal foundations in Belgium [45,80]. Assessing connectivity of colonising organisms between structures will require understanding planktonic larval ecology such as dispersive phase duration and larval behaviour, post-dispersal to adult habitat use [46] and hydrodynamics [79]. In relation to NIS the relevant scale for study could be by national sector (e.g. the Dutch North Sea) or entire regions (e.g. entire North Sea under the MSFD) depending on perspectives and spatially-defined objectives.

2.3.3. Biodiversity

Proponents of MREDs frequently cite that they will be associated with increased biodiversity [4,81]. Biodiversity is a complex, multifaceted phenomenon [82] and high biodiversity is often associated with increased system resilience to stressors ranging from physical disturbance to challenge by non-indigenous species [13,82,83]. High levels of biodiversity are, consequently, considered beneficial and actions to conserve/ enhance biodiversity are required under various international conventions (e.g. Aichi Biodiversity targets 2011–2020 under the Convention on Biological Diversity [84]). It is crucial that any assessment of biodiversity is understood to be scale-dependent: when placed in sedimentary environments the addition of hard substrata will inevitably enhance localised beta diversity (rate of change in diversity) because the structures invariably host different biological communities compared with the receiving environment. Macrobenthic alpha diversity (number of species per sample) may differ in proximity to hard-substrata mediated via structure-derived enhanced organic flux [85,86], changes in sediment stability [4] or changes in sediment attributable to colonising organisms such as mussels [87]. However, whether (how or why) increased alpha or beta diversity necessarily constitutes an enhancement to the system, on the scale of one or more ‘non-natural’ structures, has not been sufficiently explored and there is growing concern about ‘ocean sprawl’ [88,89]. There are currently no guidelines as to what levels (i.e. thresholds) of biodiversity are desirable, which assemblages (e.g. micro-, meio-, macro- and/or megabenthos) should be considered and over what scale (in space and time) it should be assessed [82]. In terms of regulatory compliance, the goal is usually to demonstrate an increasing biodiversity trend though the scales (time and space) over which this should be demonstrated and the underlying rationale are unknown [82]. Even a routine, explicit acknowledgement that appropriate scales for biodiversity-monitoring are ‘known-unknowns’ would constitute a step forward. The basis of metric choice, and thresholds, or even the acknowledgement that these cannot be determined/agreed, should be part of the environmental assessment in order to improve clarity and transparency of monitoring programmes (e.g. as required under the Marine (Scotland) Act, 2010).

2.3.4. The impact of fishing exclusion and assessing recovery

The extent to which MREDs exclude other activities (e.g. fishing) will, in part, determine their large-scale impacts on a range of receptors particularly fisheries and biodiversity [4,90]. Towed-gear fishing is banned around some operational wind-farms (e.g. in the Belgian and Dutch sectors) but not others (e.g. some in the UK sector) whilst tidal and wave devices are likely to be avoided by fishermen because of the entanglement risk. Where fishing effort exclusion occurs it may allow recovery of the benthos (with spill-over effects from increased local production extending an unknown distance beyond the MRED’s

boundary [91]). This ‘protective’ effect [81] may allow benthic recovery of reef-forming ecosystem-engineers, such as reef-forming polychaetes [92] which will have implications for local biodiversity [4] and potentially regional-scale effects if acting as a nursery ground (e.g. for fish including commercial species) or a substantial source of propagules. The appropriate spatial- and temporal -scale for monitoring the degree of benthic recovery following fishery-exclusion will depend on the objectives of the monitoring programme and the sampling effort employed. Surveys of > 10 years might be inadequate to quantify recovery of some fish species, particularly where they are depleted because of overfishing (as by-catch) [93]. Appropriate time-scales for assessing change/recovery in relation to fishery-effort cessation will vary between sites: > 10 years exclusion have resulted in a measurable difference in the North Sea benthos [94] but in other cases (e.g. a closure of 5 years) they have not [95]. There is considerable overlap between the monitoring objectives in relation to assessing the ‘efficacy’ of marine protected areas [96] and those to assess the ‘impacts’ of MREDs [6,97].

The identification of spatially/temporally delimited metrics and thresholds, in line with the overall management objectives, are the critical components to a logically based monitoring programme. However, many current monitoring programmes are deficient in this regard yet, based on null hypothesis significance testing, they still proceed [17,25](\$2.4).

2.4. Fundamental issues around data gathering and interpretation

Currently, most benthic monitoring effort focusses on detecting whether local impacts (changes) around MREDs are statistically significant or not at arbitrary spatial and temporal scales [17] and this results in DRIPy data. This approach, centred around null hypothesis significance testing (NHST), is not appropriate within the context of adaptive management as such null hypotheses are inevitably false and, in this context, failure to reject the null hypothesis merely reflects the lack of statistical power of the sampling design employed [98–100]. NHST as the mainstay of statistical analysis for the ecological/biological (and other) sciences stems from the unfortunate conflation of two different statistical approaches (Fisher v. Neyman and Pearson [100]) and has resulted in a ‘success’ being associated with a rejection of the null hypotheses. The deficiencies of NHST were acknowledged by Fisher, Neyman and Pearson in the 1950s in relation to their own fields [101] but may have persisted in scientific endeavour because it enables an attractively simple ‘statistical ritual’ to be adopted by non-statisticians (i.e. most scientists and regulators) [100]. In the context of monitoring the null-hypothesis is usually one of there being no-change⁹ in relation to a development [100]. This approach is particularly dangerous where ‘no evidence of impact’ (i.e. null hypothesis acceptance) is interpreted as ‘evidence of no impact’ [101]. There is a good argument that the obligation of proof should be reversed where environmental monitoring is concerned or, at least, that every environmental monitoring programme should fully investigate the power of the observational programme to detect threshold exceedance.

2.4.1. Null hypothesis significance testing and Type I and II errors

A common misunderstanding in relation to NHST is that an apparently simple question ‘is there an impact?’ can be unequivocally answered based around a P-value¹⁰ threshold (usually 0.05) and that,

⁹ No-change is a logical impossibility in terms of environmental monitoring because changing the environment, via MREDs, will inevitably cause change, even if not ‘significant’ from a statistical or ecological perspective.

¹⁰ A P-value can be defined as the probability of observing the data conditional on the null hypothesis being true (for a more detailed definition see [98]). It is not the probability that the null hypothesis is true as, for example, is the likely interpretation of the P-value justified statement ‘...is not significantly impacting epifaunal communities...’ (see line 1, Discussion in [135]).

consequently, a threshold does not need to be specified. This is incorrect; justified thresholds are required (§2.2.) around which to base effective management decision-making.

Accepting the logical requirement for a threshold raises two questions: first the required confidence around any assessment of threshold-exceedance that is made and, second, what should happen if the threshold is exceeded (this involves an analysis of risk, cost and liability). In environmental monitoring the meaning of the term ‘cost’ depends on perspective – a developer generally wishes to minimise the observational effort (e.g. the cost associated with taking samples) but other stakeholders will be more concerned with costs occurring as a consequence of the activity being monitored (e.g. loss of spawning habitat). Following the selection of a metric and threshold (§2.2), there needs to be agreement, between stakeholders, on the cost of either determining that the threshold is exceeded when, in reality, it is not or failing to determine threshold-exceedance when, in reality, it is occurring (known as Type I and II errors respectively). Stakeholders also need to agree how to approach decision-making in the face of varying degrees of uncertainty. In traditional monitoring programmes, the Type I error rate is set (generally, and arbitrarily, at 0.05) whilst the Type II error, which is arguably more important for conservation-risk management, is not specified and is frequently ignored [50]. A ‘total error rate’, split between Type I and II errors, has been suggested to accommodate respective costs [50,51,102]. This type of approach focuses attention on asking relevant questions (e.g. which metric to monitor) and to consider how much change is acceptable whilst balancing the stakeholder-specific costs of making both error types. Where the probabilities and costs of several outcomes are known, or can be reliably estimated, ‘decision theory’ can be used to maximise the positive outcomes [48].

2.4.2. An alternative approach to NHST

Balancing ‘precautionary’ and ‘risk-based’ decision making is core to the EIA process [103] within which monitoring is specified. Given the numerous limitations of NHST (§2.4.1) and the aspiration to manage ecosystem relevant changes, we suggest decision making be based on an upper confidence interval [6] in relation to the evidence provided by data, as illustrated in Fig. 1.

The scenario in Fig. 1 assumes that an initial threshold (defined via expert elicitation [49]) of 20 Units of Impact (UOI, Fig. 1, solid blue line) in the metric of interest has been identified beyond which change is considered unacceptable. In this example, expert opinion about the assessed impact indicates considerable uncertainty about the likely UOI (–10 to 30 UOI, Fig. 1 ‘Expert opinion’) but that it is plausible that the threshold might be exceeded (Fig. 1, ‘Risk’) and monitoring is deemed necessary. Further consultation would be required to agree the degree of confidence that should be applied to any point estimate of the relevant parameter; here that confidence is arbitrarily set at 95/99% and relates to the upper confidence boundary only (UCB, Fig. 1, horizontal solid black lines extending horizontally from means A – E). The hypothesis under test is that the threshold (20 UOI) is not exceeded; this is not a null-hypothesis of no-impact. Provided the monitoring programme is designed appropriately (not biased or subject to confounding), it will generate mean-estimates that are centred on the true UOI-parameter. The likely spread of these mean estimates, should the same sampling strategy be repeated many times, is indicated by the lower and upper confidence intervals (under the frequentist paradigm; [104]). If the actual UOI was 10, the most likely outcomes shown in Fig. 1 are A, A*, B and D, which all result in the same point estimate (~10) but with differing levels of confidence in that estimate which may result in differing management decisions. The evidence at A suggests that > 20 UOI is unlikely to be found and that monitoring could be reduced (Green). At B, the upper 95% CI indicates that 20 UOI might be exceeded and that monitoring should at least be continued (Amber) whilst the uncertainty is greatest for situation D which might, depending on circumstance, result in enhanced monitor-




Fig. 1. – ‘Traffic light’ evidence for action (reduce, maintain/enhance monitoring or enter into mitigation) as a function of the point estimate (usually the mean, indicated here by the letter) and upper 95% and 99% confidence boundary (UCB) as indicated by the solid (all) and dotted (A* only) lines. The threshold of 20 units of impact (UOI) has been agreed (by expert elicitation). Traditionally, a null hypotheses of no-impact is tested (zero UOI, double line) and this is accepted if the lower CI (Scenario G) includes it. We recommend that decisions be made on based around an agreed upper confidence interval. See main text §2.4.2 for further details. (For interpretation of the references to color in this figure, the reader is referred to the web version of this article).

ing or mitigation (Red). Entering into mitigation in these circumstances would be a precautionary decision (assuming that 10 UOI is the true value). In evaluating the risk of threshold-exceedance a balance is required between the interpretation of a point estimate and the UCB, for example, it may not be straightforward to evaluate the relative risks between B and C where B has the lower point estimate, compared with C, but the higher UCB.

The scope for Type II errors are reduced by increasing the UCB and this process can be extended indefinitely but this commensurately increases the rate of Type I errors. In the case of Fig. 1: A, had the actual impact metric been 20 UOI, a Type II error would have occurred using a 95% CI (A) but not if a 99% CI has been employed (A*) but this would not, as is typically the case, be as a consequence of under-sampling (see below).

The commonly adopted NHST approach would be to test the null hypothesis of no-change (i.e. that the UOI is zero, double blue line on Fig. 1) and make decisions based on the lower confidence interval. Fig. 1:G illustrates the results from a very weak sampling programme (large CI) and, assuming the correct value is 25 UOI (the mean of G which exceeds the threshold), one liable to abuse if used to justify inaction. The Type II error rate is proportional to sampling effort - additional sampling could result in Fig. 1:F and the correct rejection of the null even based on the lower CI. Type II errors are routinely made in relation to monitoring programmes that have insufficient power (sample size) but, conversely, where under-sampling does occur, and results in the correct rejection of a null hypothesis, then the point estimate is inevitably over-estimated because of ‘truth-inflation’ [105], Appendix A, Fig. 3). Consequently, many of the published impacts associated with MREDs, where the sample size is low, may be overestimated.

The widths of the confidence interval (precision) are synonymous with ‘statistical power’ (narrower intervals indicating greater statistical power). In a situation of limited sampling effort (budget), a balance has to be struck between precision (proportional to sampling density) against the population to which the observations can be inferred (proportional to spatial domain). How this is achieved is influenced by the characteristics of the response variable and scale of assessment




Fig. 2. – Summary of a plausible route to a rationalised monitoring programme, focussed around identified thresholds within agreed spatial and temporal domains. We recommend that, if the estimated effort/cost of a monitoring programme required to assess a change, to agreed precision, is not logistically feasible and/or cost-effective then the suggestion is not to conduct the monitoring programme. The green, amber and red colours (X, Y, Z respectively) should be cross-referenced to Fig. 1. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article).

and this should be dictated by the spatio-temporally delimited management objectives (§2.5).

Agreeing the balance between the relative risks (Type I and II errors) and confidence around parameter estimates poses major challenges in agreeing a monitoring protocol. However, justified thresholds e.g. around non-linear effects (§2.2) and pre-data-gathering agreements about how to accommodate uncertainty enable logically-based, defensible and auditable decision-making and a degree of flexibility, adaptability and compromise that is considered a part of successful EIA programmes [48,106–108].

2.5. Assessing change in time and space

With the necessary metrics, thresholds and scales identified the sampling programme must be designed. Accurately and precisely assessing change, in time and space, in highly naturally variable environments is not simple. Whilst new technologies are making the monitoring/mapping of some receptors realistic at large-scales, detect-

ing change/threshold exceedance will always be challenging in the marine environment particularly where temporal/spatial variability is high and/or poorly quantified. Field-based monitoring programmes constitute observational studies and these, inevitably, do not lend themselves to inference (assigning cause to effect) because ‘treatments’ (e.g. wind-farm monopiles) are not randomly assigned to the region of interest. In these circumstances an assessment of change over both space and time requires an understanding of pre-development spatio-temporal variability in the metric under investigation in order to conduct ‘before-after-control-impact’-(BACI)-type comparisons [109–112] and this is routinely done in relation to MRED monitoring. BACI designs are useful where the effect is manifested as an acute and/or long-lasting change in the mean of the response variable but they perform less well where the impact is gradual or manifested as a change in variability [113] as is typical of the changes likely to occur around MREDs [86]. Minimising undesirable variability (i.e. that not attributable to the impact source) in monitoring programmes will maximise their cost-effectiveness [114]. Variability can be controlled by




Fig. 3. – Example of ‘truth-inflation’. The two populations being compared have the following distributions: $X \sim N(50, 100)$ and $Y \sim N(45, 100)$ (the effect-size being estimated is therefore $50 - 45 = 5$, as indicated by the red horizontal line). When the sample size is small (here $n = 10$), in order to ‘correctly’ reject the null hypothesis ($P < 0.05$) the determined effect size (point-estimates) are at least ~ 10 (in this example run), twice the actual effect size (hence ‘truth-inflation’). Based around a P-value rejection threshold of 0.05 most of the simulations result in a ‘Type II’ error (lie to the right of the vertical green line, $P = 0.05$). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article).

conducting sampling at the same time of year [85], time of day, tidal state (if relevant) and with appropriate stratification (e.g. across different substratum types) and by including environmental and meteorological field-details in the statistical models used to make parameter predictions. Even with careful timing, naturally occurring temporal variability (e.g. as a consequence of unpredictable storm-induced resuspension events) suggests that change should be monitored over large-spatial scales [115].

Many MRED consenting authorities (e.g. in the UK, US (some States) and France) require pre-development site characterisation data to be collected, typically over a period of 1–3 years prior to development. Non-existent or poor-parameterisation of the temporal variability characterising benthic systems (i.e. as will happen when using only 1–3 years’ data) means that these approaches are only likely to detect the most extreme changes [116]. However, such comparisons have been used to support conclusions of no ‘significant’ impact with a ‘high degree of confidence’, presumably based on expert judgement of temporal variability (e.g. tidal device; Strangford Lough, UK [44],¹¹ offshore windfarm: East Anglia, UK [117]¹²). Overall, Franco et al. [14] showed that current North Sea offshore wind-farm statutory monitoring studies (UK Sector) could only reliably detect diversity/abundance changes of greater than 50%. Given the absence of a regulatory threshold or clear objectives associated with the monitoring programmes assessed by Franco et al. [14] it is impossible to say whether they were, and continue to be, fit for purpose.

Technologies, such as multibeam- and side-scan-sonar, are increas-

ingly allowing cost-effective mapping of entire MRED sites (pre- and/or post-development) and, with appropriate ground-truthing, this approach reduces the scope for sampling error because whole populations can be mapped [92,118,119]. In the absence of such technology, mapping usually consists of taking samples across the area of interest and interpolating between them. Bijleveld et al. [120] found the optimal approach to benthic mapping was to take samples from a regular-spaced grid complimented with some randomly located samples whilst Van der Meer [121], in relation to monitoring change in benthic communities over time, found that repeat sampling of randomly located stations was optimal. Data with inherent sampling dependencies (e.g. from BACI designs) pose particular statistical problems and require mixed- or generalised estimating equation (GEE) -models (see the ‘R’ package ‘MRSea’). Many monitoring programmes assess multivariate patterns using dissimilarity-matrix-based analyses [122] (e.g. Primer™ and various routines in the ‘Vegan’ library in R). These approaches are invariably adopted to test null-hypotheses of no-change (usually based on a P-value threshold of 0.05) with all the inherent problems of interpretation (§2.4). An alternative to dissimilarity-based tests is to use a multivariate model-based approach such as provided in the ‘MVabund’ R package [122] which, unlike the former, enables an assessment of which taxa are changing in relation to multiple drivers.

Understanding long-term trends (e.g. occurring over the lifetime of MREDs) requires post-development comparisons with long-term time-series data [112,123,124] and such data are relatively rare,¹³ which causes problems for benthic monitoring in relation to any marine activity. Whilst knowledge of natural variability in the populations under investigation is essential to quantify MRED-related change there is currently little guidance on what constitutes an appropriate temporal or spatial scale in relation to MRED monitoring.

3. Conclusions

The coastal zone is facing a period of unprecedented human development resulting in both new pressures and an increase in existing pressures. These pressures arise from diverse activities including the expansion of MREDs which, within the EU, are part of the ‘blue-growth’ agenda [125]. To ensure that blue-growth only occurs in harmony with the environment [6103] decision makers need relevant information [126]. The benthos is a core component of the marine ecosystem and, despite considerable monitoring effort, its ecosystem-level interaction with MREDs is poorly understood. This is because most benthic monitoring is ‘DRIPY’. In the case of MREDs, we argue that benthic monitoring programmes should consider the development within the ecosystem and in the context of ecosystem-service provision [8,19,31,127]. In this respect, the relevant scale for monitoring is likely to be relatively large, extending well outside the boundary of any particular development, and will include numerous activities and developments including multiple MREDs [31].

In order to ‘turn-off the DRIP’ we recommend the approach summarised in Fig. 2. Monitoring studies should start with clear management objectives (§2.1) in terms of relevant metrics and spatial/temporal domains (§2.2 and §2.3) and, where possible, pre-defined thresholds of unacceptable change (§2.2). The metrics, thresholds and necessary confidence in any assessment should be agreed by relevant stakeholders and consenting authorities (§2.4.2, Fig. 2: Phase I) prior to the sampling-design phase (§2.5, Fig. 2: Phase II). We

¹¹ Page 57: “Significance in this case is considered to be biological significance beyond natural variation”. Natural variation could not have been estimated from the data so this must have been based on expert judgement.

¹² Chapter 9, Page 22: “significance is determined by professional judgment ...if a predicted change is **within the range of natural variability** of the baseline environment then it is not considered to be significant”. Natural variability was not assessed as samples were only collected over a period of ~6 months.

¹³ International Bottom Trawl Survey (1970 - present; <http://ocean.ices.dk/Project/IBTS/>), the Norderney (German sector, North Sea) macrobenthic time-series data (1978 - present; [136]), the continuous plankton recorder data (1931 - present; <http://www.sahfos.ac.uk/>), the Shetland Oil Terminal macrobenthic data (1978 - present; <http://www.soteag.org.uk/>), Bay of Morlaix, western English Channel (1977- present; [137]) and from the Northumberland coast (UK, North Sea) a macrobenthic /plankton record collected from 1972 - present [138].

acknowledge that it might be difficult, given the lack of scientific understanding or precedents, to identify/agree thresholds amongst diverse stakeholder groups. Under such circumstances, we urge regulators and developers to consider the value of the data that are being generated in terms of parameterising ecosystem models (Fig. 2: Phase III) [128–130]. Where thresholds can be identified, particularly where the response metric is likely to be strongly non-linearly related to the environmental pressure (§2.2), then consideration should be given as to whether any realistic sampling programme can assess, with the necessary precision, the likelihood of the threshold being exceeded (§2.4.2, Fig. 2: Phase II). In the event that the sampling programme is not considered likely to deliver useful data then the programme should not proceed in its proposed form (i.e. ‘DRIPy’ data should be avoided) and the resources used elsewhere (Fig. 2: Phase II). Where a logical monitoring programme can be designed, regulatory decisions, including the requirement to cease, reduce, maintain or enhance monitoring/enter into mitigation, can then be fully justified to all stakeholders reducing conflict, expenditure and risk (e.g. of litigation) (Fig. 2, Phase III).

In order to understand the ecological consequences of MREDS within a rapidly growing marine sector, with multiple, potentially competing, objectives there needs to be a strategic overview of the relevant spatial domain in which monitoring could occur (§2.3) and this can be challenging to manage [31]. This strategic overview could be provided by the regulators or other competent authority and, in

Europe, this regional scale assessment could be initiated under the Strategic Environmental Assessment Directive (SEA Directive 2001/42/EC). Within Scotland (UK), the Scottish Offshore Renewables Research Framework (SpORRAn),¹⁴ established under the auspices of the Scottish Regulator (Marine Scotland) includes a benthic sub-group and is adopting a more strategic, ‘joined-up’ ecosystem-level approach to understanding MRED-benthos interactions. Funding such non-development-specific monitoring/research is challenging and we recommend the Belgian approach where developers contribute to a central funding pot from which strategic research can be funded. Supporting basic, multi-sector, ecosystem-scale research is the best way to understand, and expedite, the sustainable future of the marine renewable energy industry. We urge all concerned (developers, regulators and their advisors) to question the basis of current benthic monitoring programmes and to ‘turn-off the DRIP’.

Acknowledgements

This manuscript was developed as part of the Terms of Reference of the ICES Working Group on Marine Benthic and Renewable Energy Developments (WGMBRED) finalised during the 2015 meeting at SAMS, Oban with support from SAMS and the Marine Alliance for Science and Technology, Scotland (MASTS). TW acknowledges support from the FP7 –ENERGY project ‘TIDES’ (Grant no. 322428).

Appendix A

```
###R CODE TO DEMONSTRATE ‘TRUTH INFLATION’.
#set parameters
effectsize=5
sample.size=10
nos.iterations=20
SDev=10
Mean=50
Pvalues=NULL; EffectSizeEstimate=NULL; StandardError=NULL
TestData=function(){
d=data.frame(Treatment = gl(2, k=sample.size, length=2*sample.size, labels = c('Low', 'High')),
  Test = c(rnorm(n=sample.size, mean=Mean, sd=SDev),
    rnorm(n=sample.size, mean=Mean-effectsize, sd=SDev)))
  return(d)}
lmp <- function (modelobject) {
  if (class(modelobject) != "lm") stop("Not an object of class 'lm' ")
  f <- summary(modelobject)$fstatistic
  p <- pf(f [1],f [2],f [3], lower.tail=F)
  attributes(p) <- NULL
  return(p)
}
for(i in 1: nos.iterations){
d=TestData()
M1=lm(Test~Treatment, data=d);
Pvalues[i]=(lmp(M1))
EffectSizeEstimate[i]= mean(d$Test[d$Treatment=="Low"])-mean(d$Test[d$Treatment=="High"])
StandardError[i]=summary(M1)$coefficients[2,2]
}
TestDataSet=data.frame(cbind(Pvalues, EffectSizeEstimate, StandardError))#
require(ggplot2)
Pvalueplot=ggplot(TestDataSet, aes(x = Pvalues, y = EffectSizeEstimate)) +
geom_point(size = 4) +
geom_errorbar(aes(ymin=EffectSizeEstimate-qt(0.975, sample.size)*StandardError,
  ymax=EffectSizeEstimate+qt(0.975, sample.size)*StandardError))+
geom_hline (yintercept=effectsize, col="red", lwd=1.5)+
```

¹⁴ <http://www.gov.scot/Topics/marine/Licensing/marine/scoping/orelg/SpORRAn>

```

geom_vline (xintercept=0.05, col="green", lwd=1.5)+
labs(title="Simulated data, true effect size as indicated by red line")+
labs(x="Determined P value")+
labs(y="Effect size estimate")+
theme_bw()
Pvalueplot
P2=Pvalueplot + theme(axis.title = element_text(size=20),
axis.text = element_text(size=20))
P2
P2+ annotate("text", x=c(0.76,0.8,0.76), y=c(20,21,22),
label=c(paste("Sample size=", sample.size),
paste("SDev=", SDev),
paste("Effect size=", (effectsize/Mean)*100, "%")))
MeanEffectEstimate=mean(TestDataSet$EffectSizeEstimate[TestDataSet$Pvalue <=0.05])
print(paste("The actual effect size =", effectsize,
"while the mean determined effect size when P <=0.05 is", signif(MeanEffectEstimate,3)))

```

References

- Costanza R, de Groot R, Sutton P, van der Ploeg S, Anderson SJ, Kubiszewski I, et al. Changes in the global value of ecosystem services. *Glob Environ Chang* 2014;26:152–8. <http://dx.doi.org/10.1016/j.gloenvcha.2014.04.002>.
- Miller RG, Hutchison ZL, Macleod AK, Burrows MT, Cook EJ, Last KS, et al. Marine renewable energy development: assessing the Benthic Footprint at multiple scales. *Front Ecol Environ* 2013;11:433–40. <http://dx.doi.org/10.1890/120089>.
- Boehlert GW, Gill AB. Environmental and ecological effects of ocean renewable energy development: a current synthesis. *Oceanography* 2010;23:68–81.
- Inger R, Attrill MJ, Bearhop S, Broderick AC, Grecian WJ, Hodgson DJ, et al. Marine renewable energy: potential benefits to biodiversity? An urgent call for research. *J Appl Ecol* 2009;46:1145–53. <http://dx.doi.org/10.1111/j.1365-2664.2009.01697.x>.
- Borger T, Hattam C, Burdon D, Atkins JP, Austen MC. Valuing conservation benefits of an offshore marine protected area. *Ecol Econ* 2014;108:229–41. <http://dx.doi.org/10.1016/j.ecolecon.2014.10.006>.
- Maclean IMD, Inger R, Benson D, Booth CG, Embling CB, Grecian WJ, et al. Resolving issues with environmental impact assessment of marine renewable energy installations. *Front Mar Sci* 2014;1:1–5. <http://dx.doi.org/10.3389/fmars.2014.00075>.
- Judd AD, Backhaus T, Goodsir F. An effective set of principles for practical implementation of marine cumulative effects assessment. *Environ Sci Policy* 2015;54:254–62. <http://dx.doi.org/10.1016/j.envsci.2015.07.008>.
- O'Higgins T. A systems approach to environmental management. Dunedin Academic Press; 2015.
- Galparsoro I, Borja A, Uyarra MC. Mapping ecosystem services provided by benthic habitats in the European North Atlantic Ocean. *Front Mar Sci* 2014. <http://dx.doi.org/10.3389/fmars.2014.00023>.
- Hewitt JE, Ellis JI, Thrush SF. Multiple stressors, nonlinear effects and the implications of climate change impacts on marine coastal ecosystems. *Glob Chang Biol* 2015. <http://dx.doi.org/10.1111/gcb.13176>.
- Hunsicker M, Kappel C, Selkoe KA, Halpern BS. Characterizing driver-response relationships in marine pelagic ecosystems for improved ocean management. *Ecol Appl* 2015. <http://dx.doi.org/10.1007/s13398-014-0173-7>.
- Mee LD, Jefferson RL, Laffoley D da, Elliott M. How good is good? Human values and Europe's proposed marine strategy directive. *Mar Pollut Bull* 2008;56:187–204. <http://dx.doi.org/10.1016/j.marpolbul.2007.09.038>.
- Tett P, Gowen RJ, Painting SJ, Elliott M, Forster R, Mills DK, et al. Framework for understanding marine ecosystem health. *Mar Ecol Prog Ser* 2013;494:1–27. <http://dx.doi.org/10.3354/meps10539>.
- Franco A, Quintino V, Elliott M. Benthic monitoring and sampling design and effort to detect spatial changes: a case study using data from offshore wind farm sites. *Ecol Indic* 2015;57:298–304. <http://dx.doi.org/10.1016/j.ecoind.2015.04.040>.
- Reynolds P, Vella G, Freeman S, May J, Savage A, Jenner C. Consenting lessons learned - an offshore wind industry review of past concerns, lessons learned and future challenges. *RenewablesUK* 2011.
- Ward R, Loftis J, McBride G. The "data-rich but information-poor" syndrome in water quality monitoring. *Environ Manag* 1986;10:291–7. <http://dx.doi.org/10.1007/BF01867251>.
- MMO. Review of Post-Consent Offshore Wind Farm Monitoring Data Associated with Licence Conditions. A report produced for the Marine Management Organisation. MMO Project No: 1031; 2014, p. 194.
- Leeney RH, Greaves D, Conley D, O'Hagan AM. Environmental Impact Assessments for wave energy developments - Learning from existing activities and informing future research priorities. *Ocean Coast Manag* 2014;99:14–22. <http://dx.doi.org/10.1016/j.ocecoaman.2014.05.025>.
- Papathanasopoulou E, Beaumont N, Hooper T, Nunes J, Queiros AM. Energy systems and their impacts on marine ecosystem services. *Renew Sustain Energy Rev* 2015;52:917–26. <http://dx.doi.org/10.1016/j.rser.2015.07.150>.
- Villnäs A, Norkko A. Benthic diversity gradients and shifting baselines: implications for assessing environmental status. *Ecol Appl* 2011;21:2172–86. <http://dx.doi.org/10.1890/1014-7331>.
- Arntz WE, Gili JM, Reise K. Unjustifiably Ignored: reflections on the role of benthos in marine ecosystems. In: Gray JS, Ambrose W, Szaniawska A, editors. *Biogeochemical cycling sediment ecology*, Dordrecht, Netherlands: Springer; 1999. p. 105–24. http://dx.doi.org/10.1007/978-94-011-4649-4_5.
- Schmitt RJ, Osenberg CW. Detecting ecological impacts caused by human activities. In: Schmitt RJ, Osenberg CW, editors. *Detecting ecological impacts*. San Diego: Academic Press; 1996. p. 3–15.
- Lindeboom H, Degraer S, Dannheim J, Gill AB, Wilhelmsson D. Offshore wind park monitoring programmes, lessons learned and recommendations for the future. *Hydrobiologia* 2015;169–80. <http://dx.doi.org/10.1007/s10750-015-2267-4>.
- Kannen A. Challenges for marine spatial planning in the context of multiple sea uses, policy arenas and actors based on experiences from the German North Sea. *Reg Environ Chang* 2014;14:2139–50. <http://dx.doi.org/10.1007/s10113-012-0349-7>.
- Maclean IMD. Resolving issues with environmental impact assessment of marine renewable energy installations. *Front Mar Sci* 2014;1:1–5. <http://dx.doi.org/10.3389/fmars.2014.00075>.
- Blenckner T, Kannen A, Barausse A, Fischer C, Heymans JJ, Luisetti T. Past and future challenges in managing European seas. *Ecol Soc* 2015;20:40. <http://dx.doi.org/10.5751/es-07246-200140>.
- Freire-Gibb LC, Koss R, Margonski P, Papadopoulou N. Governance strengths and weaknesses to implement the marine strategy framework directive in European waters. *Mar Policy* 2014;44:172–8. <http://dx.doi.org/10.1016/j.marpol.2013.08.025>.
- Bergström L, Kautsky L, Malm T, Rosenberg R, Wahlberg M, Åstrand Capetillo N, et al. Effects of offshore wind farms on marine wildlife—a generalized impact assessment. *Environ Res Lett* 2014;9:1–12. <http://dx.doi.org/10.1088/1748-9326/9/3/034012>.
- Willsted E, Gill AB, Birchenough SNR, Jude S. Assessing the cumulative environmental effects of marine renewable energy developments: establishing common ground. *Sci Total Environ* 2017;577:19–32. <http://dx.doi.org/10.1016/j.scitotenv.2016.10.152>.
- Brennan J, Fitzsimmons C, Gray T, Raggatt L. EU marine strategy framework directive (MSFD) and marine spatial planning (MSP): which is the more dominant and practicable contributor to maritime policy in the UK?. *Mar Policy* 2014;43:359–66. <http://dx.doi.org/10.1016/j.marpol.2013.07.011>.
- Barrio Froján CRS, Cooper KM, Bolam SG. Towards an integrated approach to marine benthic monitoring. *Mar Pollut Bull* 2016;104:20–8. <http://dx.doi.org/10.1016/j.marpolbul.2016.01.054>.
- Bailey H, Brookes KL, Thompson PM. Assessing environmental impacts of offshore wind farms: lessons learned and recommendations for the future. *Aquat Biosci* 2014;10:8. <http://dx.doi.org/10.1186/2046-9063-10-8>.
- Boyd I, Bowen D, Iverson S. *Marine mammal ecology and conservation*. Oxford, UK: Oxford University Press; 2010.
- JNCC. Seabird Monitoring Programme. (<http://jncc.defra.gov.uk/page-1550>) [Accessed Mar 2016].
- Russell DJF, SMJM Brasseur, Thompson D, Hastie GD, Janik VM, Aarts G, et al. Marine mammals trace anthropogenic structures at sea. *Curr Biol* 2014;24:R638–R639. <http://dx.doi.org/10.1016/j.cub.2014.06.033>.
- Fox CJ, Benjamins S, Masden E, Miller R. Challenges and opportunities in monitoring the impacts of tidal-stream energy devices on marine vertebrates. *Renew Sustain Energy Rev* 2017, [In press].
- King SL, Schick RS, Donovan C, Booth CG, Burgman M, Thomas L, et al. An interim framework for assessing the population consequences of disturbance. *Methods Ecol Evol* 2015;6:1150–8. <http://dx.doi.org/10.1111/2041-210X.12411>.

- [38] Akçakaya H, Ferson S, Burgman MA, Keith DA, Mace GM, Todd CR. Making consistent IUCN classifications under uncertainty. *Conserv Biol* 2000;14:1001–13. <http://dx.doi.org/10.1046/j.1523-1739.2000.99125.x>.
- [39] Walters CJ, Martell SJD. Fisheries ecology and management. New Jersey, US: Princeton University Press; 2004. <http://dx.doi.org/10.1046/j.1365-2400.1999.00126.x>.
- [40] Large SI, Fay G, Friedland KD, Link JS. Defining trends and thresholds in responses of ecological indicators to fishing and environmental pressures. *ICES J Mar Sci* 2013;70:755–67.
- [41] MMO. Review of Post-Consent Offshore Wind Farm Monitoring Data Associated with Licence Conditions. A report produced for the Marine Management Organisation. MMO Project No: 1031. ISBN 978-1-909452-24-4; 2014, p. 194.
- [42] Aumüller R, Baier H, Binder A, Damian H, Feindt-herr H, Merck T, et al. Standard Investigation of the Impacts of offshore wind turbines on the marine environment (StUK4); 2013. <http://www.bsh.de/en/Products/Books/Standard/7003eng.pdf> [Accessed Mar 2017].
- [43] Sheehan EV, Stevens TF, Attrill MJ. A quantitative, non-destructive methodology for habitat characterisation and benthic monitoring at offshore renewable energy developments. *PLoS One* 2010;5. <http://dx.doi.org/10.1371/journal.pone.0014461>.
- [44] Keenan G, Sparling C, Williams H, Fortune F. SeaGen Environmental Monitoring Programme. (<http://seagenation.co.uk/files/SeaGen-Environmental-Monitoring-Programme-Final-Report.pdf>) [Accessed Mar 2016 2011].
- [45] De Mesel I, Kerckhof F, Norro A, Rumes B, Degraer S. Succession and seasonal dynamics of the epifauna community on offshore wind farm foundations and their role as stepping stones for non-indigenous species. *Hydrobiologia* 2015;756:37–50. <http://dx.doi.org/10.1007/s10750-014-2157-1>.
- [46] Reubens JF, Degraer S, Vincx M. The ecology of benthopelagic fishes at offshore wind farms: a synthesis of 4 years of research. *Hydrobiologia* 2014;727:121–36. <http://dx.doi.org/10.1007/s10750-013-1793-1>.
- [47] Hattam C, Atkins JP, Beaumont N, Börger T, Böhnke-henrichs A, Burdon D, et al. Marine ecosystem services: linking indicators to their classification. *Ecol Indic* 2015;49:61–75. <http://dx.doi.org/10.1016/j.ecolind.2014.09.026>.
- [48] Polasky S, Carpenter SR, Folke C, Keeler B. Decision-making under great uncertainty: environmental management in an era of global change. *Trends Ecol Evol* 2011;26:398–404. <http://dx.doi.org/10.1016/j.tree.2011.04.007>.
- [49] Addison PFE, de Bie K, Rumpff L. Setting conservation management thresholds using a novel participatory modeling approach. *Conserv Biol* 2015;29:1411–22. <http://dx.doi.org/10.1111/cobi.12544>.
- [50] Mapstone BD. Scalable decision rules for environmental impact studies: effect size, type I, and type II errors. *Ecol Appl* 1995;5:401–10. <http://dx.doi.org/10.2307/1942031>.
- [51] Mudge JF, Baker LF, Edge CB, Houlihan JE. Setting an optimal alpha that minimizes errors in null hypothesis significance tests. *PLoS One* 2012;7. <http://dx.doi.org/10.1371/journal.pone.0032734>.
- [52] Borja A, Miles A, Ochoipinti-Ambrogi A, Berg T. Current status of macroinvertebrate methods used for assessing the quality of European marine waters: implementing the Water framework directive. *Hydrobiologia* 2009;633:181–96. <http://dx.doi.org/10.1007/s10750-009-9881-y>.
- [53] Borja A, Prins TC, Simboura N, Andersen JH, Berg T, Marques J-C, et al. Tales from a thousand and one ways to integrate marine ecosystem components when assessing the environmental status. *Front Mar Sci* 2014;1:1–20. <http://dx.doi.org/10.3389/fmars.2014.00072>.
- [54] WMGM van Loon, Boon, Gittenberger AR, Walvoort A, Lavaley DJJ, Duineveld M. GCA, et al. application of the benthic ecosystem quality index to benthos in Dutch transitional and coastal waters. *J Sea Res* 2015;103:1–13. <http://dx.doi.org/10.1016/j.seares.2015.05.002>.
- [55] Dauvin JC, Andrade H, De-La-Ossa-Carretero JA, Del-Pilar-Ruso Y, Riera R. Polychaete/amphipod ratios: an approach to validating simple benthic indicators. *Ecol Indic* 2016;63:89–99. <http://dx.doi.org/10.1016/j.ecolind.2015.11.055>.
- [56] Baird DJ, Van den Brink PJ, Chariton AA, Dafforn KA, Jahnston EL. New diagnostics for multiply stressed marine and freshwater ecosystems: integrating models, ecoinformatics and big data. *Mar Freshw Res* 2016;67:391–2.
- [57] Vennell R, Funke SW, Draper S, Stevens C, Divett T. Designing large arrays of tidal turbines: a synthesis and review. *Renew Sustain Energy Rev* 2015;41:454–72. <http://dx.doi.org/10.1016/j.rser.2014.08.022>.
- [58] Dafforn KA, Johnston EL, Ferguson A, Humphrey CL, Monk W, Nichols SJ, et al. Big data opportunities and challenges for assessing multiple stressors across scales in aquatic ecosystems. *Mar Freshw Res* 2016;67:393–413.
- [59] Black KD, Cromeey CJ. The scientific basis of marine fish farm regulation. *Sci Diliman* 2008;20:11–22.
- [60] CMACS. Benthic Survey 2010 Technical Report. CMACS Ref: J3151 Beatrice Benthic Technical Report; 2011. (<http://sse.com/media/341142/ES-Volume-4-Annex-10A-Benthic-Ecology-Technical-Report.pdf>).
- [61] Long ER, Dutch M, Partridge V, Weakland S, Welch K. Revision of sediment quality triad indicators in Puget Sound (Washington, USA): I. A Sediment Chemistry Index and targets for mixtures of toxicants. *Integr Environ Assess Manag* 2013;9:31–49. <http://dx.doi.org/10.1002/ieam.1309>.
- [62] Wilson JC, Elliott M. The habitat-creation potential of offshore wind farms. *Wind Energy* 2009;12:203–12. <http://dx.doi.org/10.1002/we.324>.
- [63] Furness RW, Wade HM, Robbins AMC, Masden EA. From tidal stream turbines and wave energy devices. *ICES J Mar Sci* 2012;69:1466–79.
- [64] JNCC UK. Regional Seas. Peterborough, England, UK 2014. (<http://jncc.defra.gov.uk/page-1612>) [Accessed May 2016].
- [65] UNESCO. Marine Spatial Planning Initiative. (<http://www.unesco-loc-marinesp.be/>) [Accessed May 2016].
- [66] Degraer S, Brabant R, Rumes R. Environmental impacts of offshore windfarms in the Belgian part of the North Sea – learning from the past to optimise future monitoring programmes; Royal Belgium Institute of Natural Sciences; 2013. (<http://www.vliz.be/en/imis?Module=ref&refid=231864&printversion=1&dropIMIStile=1>).
- [67] Asmus H, Asmus RM. The Comparative Roles of Suspension-Feeders in Ecosystems: Proceedings of the NATO Advanced Research Workshop on The Comparative Roles of Suspension-Feeders in Ecosystems Nida, Lithuania 4–9 October 2003. In: Dame RF, Olenin S, editors., Dordrecht: Springer Netherlands; 2005, p. 199–219. doi: (http://doi.org/10.1007/1-4020-3030-4_12).
- [68] Grant J, Bugden G, Horne E, Archambault M-C, Carreau M. Remote sensing of particle depletion by coastal suspension-feeders. *Can J Fish Aquat Sci* 2007;64:387–90. <http://dx.doi.org/10.1139/f07-021>.
- [69] Mckindsey CW, Archambault P, Callier MD, Olivier F. Influence of suspended and off-bottom mussel culture on the sea bottom and benthic habitats: a review. *Can J Zool* 2011;89:622–46. <http://dx.doi.org/10.1139/z11-037>.
- [70] Inglis GJ, Gust N. Potential indirect effects of shellfish culture on the reproductive success of benthic predators. *J Appl Ecol* 2003;40:1077–89. <http://dx.doi.org/10.1111/j.1365-2664.2003.00860.x>.
- [71] Joschko TJ, Buck BH, Gutow L, Schröder A. Colonization of an artificial hard substrate by *Mytilus edulis* in the German Bight. *Mar Biol Res* 2008;4:350–60. <http://dx.doi.org/10.1080/1745100801947043>.
- [72] Janßen H, Augustin CB, Hinrichsen HH, Kube S. Impact of secondary hard substrate on the distribution and abundance of *Aurelia aurita* in the western Baltic Sea. *Mar Pollut Bull* 2013;75:224–34. <http://dx.doi.org/10.1016/j.marpolbul.2013.07.027>.
- [73] Lo WT, Purcell JE, Hung JJ, Su HM, Hsu PK. Enhancement of jellyfish (*Aurelia aurita*) populations by extensive aquaculture rafts in a coastal lagoon in Taiwan. *ICES J Mar Sci* 2009;65:453–61.
- [74] Villareal TA, Hanson S, Quaila S, Jester ELE, Granade HR, Dickey RW. Petroleum production platforms as sites for the expansion of ciguatera in the northwestern Gulf of Mexico. *Harmful Algae* 2007;6:253–9. <http://dx.doi.org/10.1016/j.hal.2006.08.008>.
- [75] Chariton AA, Sun M, Gibson J, Webb JA, Leung KMY, Hickey CW, et al. Emergent technologies and analytical approaches for understanding the effects of multiple stressors in aquatic environments. *Mar Freshw Res* 2016;67:414–28.
- [76] Baxter EJ, Rodger HD, McAllen R, Doyle TK. Gill disorders in marine-farmed salmon: investigating the role of hydrozoan jellyfish. *Aquac Environ Interact* 2011;1:245–57. <http://dx.doi.org/10.3354/aei00024>.
- [77] Purcell JE, Uye SI, Lo WT. Anthropogenic causes of jellyfish blooms and their direct consequences for humans: a review. *Mar Ecol Prog Ser* 2007;350:153–74. <http://dx.doi.org/10.3354/meps07093>.
- [78] Thorpe SA. On the biological connectivity of oil and gas platforms in the North Sea. *Mar Pollut Bull* 2012;64:2770–81. <http://dx.doi.org/10.1016/j.marpolbul.2012.09.011>.
- [79] Adams TP, Miller RG, Aleynik D, Burrows MT. Offshore marine renewable energy devices as stepping stones across biogeographical boundaries. *J Appl Ecol* 2015;51:330–8. <http://dx.doi.org/10.1111/1365-2664.12207>.
- [80] Kerckhof F, Degraer S, Norro A, Rumes B. Offshore intertidal hard substrata: a new habitat promoting non-indigenous species in the Southern North Sea: an exploratory study. *R Belg Inst Nat Sci* 2011;32:27–37.
- [81] Tononen HM, Lindeboom HJ. Dark green electricity comes from the sea: capitalizing on ecological merits of offshore wind power?. *Renew Sustain Energy Rev* 2015;42:1023–33. <http://dx.doi.org/10.1016/j.rser.2014.10.043>.
- [82] Laurila-Pant M, Lehtikoinen A, Uusitalo L, Venesjärvi R. How to value biodiversity in environmental management?. *Ecol Indic* 2015;55:1–11. <http://dx.doi.org/10.1016/j.ecolind.2015.02.034>.
- [83] Díaz S, Fargione J, Chapin IIFS, I, Tilman D. Biodiversity Loss Threatens Human well-being. *PLoS Biol* 2006;4:e277. <http://dx.doi.org/10.1371/journal.pbio.0040277>.
- [84] Beaumont NJ, Austen MC, Atkins JP, Burdon D, Degraer S, Dentinho TP, et al. Identification, definition and quantification of goods and services provided by marine biodiversity: implications for the ecosystem approach. *Mar Pollut Bull* 2007;54:253–65. <http://dx.doi.org/10.1016/j.marpolbul.2006.12.003>.
- [85] Wilding TA. Effects of man-made structures on sedimentary oxygenation: extent, seasonality and implications for offshore renewables. *Mar Environ Res* 2014;97:39–47. <http://dx.doi.org/10.1016/j.marenvres.2014.01.011>.
- [86] Coates DA, Deschutter Y, Vincx M, Vanaverbeke J. Enrichment and shifts in macrobenthic assemblages in an offshore wind farm area in the Belgian part of the North Sea. *Mar Environ Res* 2014;95:1–12. <http://dx.doi.org/10.1016/j.marenvres.2013.12.008>.
- [87] Wilding TA, Nickell TD. Changes in benthos associated with mussel (*Mytilus edulis* L.) farms on the west-coast of Scotland. *PLoS One* 2013;8. <http://dx.doi.org/10.1371/journal.pone.0068313>.
- [88] Firth LB, Knights AM, Bridger D, Evans AJ, Mieszowska N, Moore PJ, et al. Ocean sprawl: challenges and opportunities for biodiversity management in a changing world. *Annu Rev* 2016;54:193–269.
- [89] Heery EC, Bishop MJ, Bugnot LCAB, Airoidi L, Mayer-Pinto M, Sheehan E V., et al. Identifying the consequences of ocean sprawl for sedimentary habitats. *J Exp Mar Bio Ecol* 2017 .
- [90] Moland E, Olsen EM, Knutsen H, Garrigou P, Espeland SH, Kleiven AR, et al. Lobster and cod benefit from small-scale northern marine protected areas: inference from an empirical before-after control-impact study. *Proc R Soc B* 2013;280:1–9. <http://dx.doi.org/10.1098/rspb.2012.2679>.
- [91] Lindeboom H, Degraer S, Dannheim J, Gill AB, Wilhelmsson D. Offshore wind park monitoring programmes, lessons learned and recommendations for the

- future. *Hydrobiologia* 2015;756:169–80. <http://dx.doi.org/10.1007/s10750-015-2267-4>.
- [92] Pearce B, Fariñas-Franco JM, Wilson C, Pitts J, deBurgh A, Somerfield PJ. Repeated mapping of reefs constructed by *Sabellaria spinulosa* Leuckart 1849 at an offshore wind farm site. *Cont Shelf Res* 2014;83:3–13. <http://dx.doi.org/10.1016/j.csr.2014.02.003>.
- [93] Maxwell D, Jennings S. Power of monitoring programmes to detect decline and recovery of rare and vulnerable fish. *J Appl Ecol* 2005;42:25–37. <http://dx.doi.org/10.1111/j.1365-2664.2005.01000.x>.
- [94] Duineveld GCA, Bergman MJN, Lavaleye MSS. Effects of an area closed to fisheries on the composition of the benthic fauna in the southern North Sea. *ICES J Mar Sci* 2007;64:899–908. <http://dx.doi.org/10.1093/icesjms/fsm029>.
- [95] Bergman MJN, Ubels SM, Duineveld GCA, Meesters EWG. Effects of a 5-year trawling ban on the local benthic community in a wind farm in the Dutch coastal zone. *ICES J Mar Sci J Du Cons* 2014. <http://dx.doi.org/10.1093/icesjms/fsu193>.
- [96] Moland E, Olsen EM, Knutsen H, Garrigou P, Espeland SH, Kleiven AR, et al. Lobster and cod benefit from small-scale northern marine protected areas: inference from an empirical before-after control-impact study. *Proc R Soc B* 2013;280:1–9. <http://dx.doi.org/10.1098/rspb.2012.2679>.
- [97] Halpern BS, Lester SE, McLeod KL. Placing marine protected areas onto the ecosystem-based management seascape. *Proc Natl Acad Sci USA* 2010;107:18312–7. <http://dx.doi.org/10.1073/pnas.0908503107>.
- [98] Goodman S. A dirty dozen: twelve P-value misconceptions. *Semin Hematol* 2008;45:135–40. <http://dx.doi.org/10.1053/j.seminhematol.2008.04.003>.
- [99] Johnson DH. The Insignificance of Statistical significance testing. *J Wildl Manag* 1999;63:763–72. <http://dx.doi.org/10.2307/3802789>.
- [100] Gigerenzer G. Mindless statistics. *J Socio Econ* 2004;33:587–606.
- [101] Stewart-Oaten A. Goals in environmental monitoring. In: Schmitt RJ, Osenberg CW, editors. *Detecting Ecological Impacts*. San Diego: Academic Press; 1996. p. 17–28.
- [102] Field SA, Tyre AJ, Jonzén N, Rhodes JR, Possingham HP. Minimizing the cost of environmental management decisions by optimizing statistical thresholds. *Ecol Lett* 2004;7:669–75. <http://dx.doi.org/10.1111/j.1461-0248.2004.00625.x>.
- [103] Wright G. Strengthening the role of science in marine governance through environmental impact assessment: a case study of the marine renewable energy industry. *Ocean Coast Manag* 2014;99:23–30. <http://dx.doi.org/10.1016/j.ocecoaman.2014.07.004>.
- [104] Nakagawa S, Cuthill IC. Effect size, confidence interval and statistical significance: a practical guide for biologists. *Biol Rev* 2007;82:591–605. <http://dx.doi.org/10.1111/j.1469-185X.2007.00027.x>.
- [105] Reinhart A. *Statistics done wrong: the woefully complete guide*. San Francisco, California, US: No Starch Press, Inc; 2015.
- [106] Lawrence DP. Impact significance determination—designing an approach. *Environ Impact Assess Rev* 2007;27:730–54. <http://dx.doi.org/10.1016/j.eiar.2007.02.012>.
- [107] Lawrence DP. Impact significance determination-back to basics. *Environ Impact Assess Rev* 2007;27:755–69. <http://dx.doi.org/10.1016/j.eiar.2007.02.011>.
- [108] Lawrence DP. Impact significance determination-pushing the boundaries. *Environ Impact Assess Rev* 2007;27:770–88. <http://dx.doi.org/10.1016/j.eiar.2007.02.010>.
- [109] Underwood AJ. On beyond BACI: sampling designs that might reliably detect environmental disturbances. *Ecol Appl* 1994;4:3–15. <http://dx.doi.org/10.2307/1942110>.
- [110] Underwood AJ. Beyond BACI - the detection of environmental impacts on populations in the real, but variable world. *J Exp Mar Biol Ecol* 1992;161:145–78.
- [111] Underwood AJ. Beyond BACI – experimental designs for detecting human environmental impacts on temporal variations in natural populations. *Aust J Mar Freshw Res* 1991;42:569–87.
- [112] Magurran AE, Baillie SR, Buckland ST, Dick JM, Elston DA, Scott EM, et al. Long-term datasets in biodiversity research and monitoring: assessing change in ecological communities through time. *Trends Ecol Evol* 2010;25:574–82. <http://dx.doi.org/10.1016/j.tree.2010.06.016>.
- [113] Underwood AJ, Chapman MG. Power, precaution, type II error and sampling design in assessment of environmental impacts. *J Exp Mar Biol Ecol* 2003;296:49–70. [http://dx.doi.org/10.1016/S0022-0981\(03\)00304-6](http://dx.doi.org/10.1016/S0022-0981(03)00304-6).
- [114] Degraer S, Brabant R, De Mesel I. Reloading basic environmental monitoring of offshore wind farms in Belgium: Phase II. In: Degraer S, Brabant R, Rumes B, Vigin L, editors. *Environmental impacts of offshore Wind Farms*. Belgian part North Sea. *Environ. impact Monit*. 2016, p. 1–16.
- [115] Armonies W. On the spatial scale needed for benthos community monitoring in the coastal North Sea. *J Sea Res* 2000;43:121–33. [http://dx.doi.org/10.1016/S1385-1101\(00\)00008-3](http://dx.doi.org/10.1016/S1385-1101(00)00008-3).
- [116] Wilding TA, Nickell TD, Hughes DJ, Narayanaswamy BE, Burrows M, Hausrath J, et al. Statistical advice to the marine habitats monitoring project under framework agreement C10-206-0387. Joint Nature Conservation Committee 2014.
- [117] SPR. East Anglia ONE Offshore windfarm - Environmental Statement, Volume 2, Offshore. Chapter 9 – Benthic and Epibenthic Environment (including Shellfish); 2012. (<https://infrastructure.planninginspectorate.gov.uk/wp-content/uploads/projects/EN010025/EN010025-000>). .
- [118] Strong JA, Elliott M. The value of remote sensing techniques in supporting effective extrapolation across multiple marine spatial scales. *Mar Pollut Bull* 2017. <http://dx.doi.org/10.1016/j.marpolbul.2017.01.028>.
- [119] JNCC. Marine Monitoring Method Finder; 2016. (<http://jncc.defra.gov.uk/page-7171>).
- [120] Bijleveld AI, van Gils JA, van der Meer J, Dekinga A, Kraan C, van der Veer HW, et al. Designing a benthic monitoring programme with multiple conflicting objectives. *Methods Ecol Evol* 2012;3:526–36. <http://dx.doi.org/10.1111/j.2041-210X.2012.00192.x>.
- [121] Van der Meer J. Sampling design of monitoring programmes for marine benthos: a comparison between the use of fixed versus randomly selected stations. *J Sea Res* 1997;37:167–79. [http://dx.doi.org/10.1016/S1385-1101\(97\)00007-5](http://dx.doi.org/10.1016/S1385-1101(97)00007-5).
- [122] Wang Y, Naumann U, Wright ST, Warton DI. Mvabund- an R package for model-based analysis of multivariate abundance data. *Methods Ecol Evol* 2012;3:471–4. <http://dx.doi.org/10.1111/j.2041-210X.2012.00190.x>.
- [123] Addison PFE, Flander LB, Cook CN. Are we missing the boat? Current uses of long-term biological monitoring data in the evaluation and management of marine protected areas. *J Environ Manag* 2015;149:148–56. <http://dx.doi.org/10.1016/j.jenvman.2014.10.023>.
- [124] Charton JAG, Williams ID, Ruzafa AP, Milazzo M, Chemello R, Marcos C, et al. Evaluating the ecological effects of Mediterranean marine protected areas: habitat, scale and the natural variability of ecosystems. *Environ Conserv* 2000;27:159–78. <http://dx.doi.org/10.1017/S037689290000199>.
- [125] EC. Communication from The Commission to the European Parliament, The Council, The European Economic and Social Committee and The Committee of the Regions: Innovation in the Blue Economy: realising the potential of our seas and oceans for jobs and growth; 2014.
- [126] Boyd I. A standard for policy-relevant science. *Nature* 2013;501:159–60.
- [127] Hooper T, Beaumont N, Hattam C. The implications of energy systems for ecosystem services: a detailed case study of offshore wind. *Renew Sustain Energy Rev* 2017;70:230–41. <http://dx.doi.org/10.1016/j.rser.2016.11.248>.
- [128] Raoux A, Tecchio S, Pezy JP, Lassalle G, Degraer S, Wilhelmsson D, et al. Benthic and fish aggregation inside an offshore wind farm: which effects on the trophic web functioning?. *Ecol Indic* 2017;72:33–46. <http://dx.doi.org/10.1016/j.ecoind.2016.07.037>.
- [129] van der Molen J, Smith HCM, Lepper P, Limpenny S, Rees J. Predicting the large-scale consequences of offshore wind turbine array development on a North Sea ecosystem. *Cont Shelf Res* 2014;85:60–72. <http://dx.doi.org/10.1016/j.csr.2014.05.018>.
- [130] Hyder K, Rossberg AG, Allen JI, Austen MC, Barciela RM, Bannister HJ, et al. Making modelling count - increasing the contribution of shelf-seas community and ecosystem models to policy development and management. *Mar Policy* 2015;61:291–302. <http://dx.doi.org/10.1016/j.marpol.2015.07.015>.
- [131] Pauly D. Anecdotes and the shifting baseline syndrome of fisheries. *Trends Ecol Evol* 1995;10:430.
- [132] Thrush ASF, Hewitt JE, Cummings VJ, Dayton PK, Cryer M, Turner SJ, et al. Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecol Appl* 1998;8:866–79.
- [133] Thurstan RH, Hawkins JP, Raby L, Roberts CM. Oyster (*Ostrea edulis*) extirpation and ecosystem transformation in the firth of Forth, Scotland. *J Nat Conserv* 2013;21:253–61. <http://dx.doi.org/10.1016/j.jnc.2013.01.004>.
- [134] Borja A, Dauer DM, Gremare A. The importance of setting targets and reference conditions in assessing marine ecosystem quality. *Ecol Indic* 2012;12:1–7. <http://dx.doi.org/10.1016/j.ecoind.2011.06.018>.
- [135] O'Carroll JPP, Kennedy RM, Savidge G. Identifying relevant scales of variability for monitoring epifaunal reef communities at a tidal energy extraction site. *Ecol Indic* 2017;73:388–97. <http://dx.doi.org/10.1016/j.ecoind.2016.10.005>.
- [136] Kröncke I, Reiss H. Influence of macrofauna long-term natural variability on benthic indices used in ecological quality assessment. *Mar Pollut Bull* 2010;60:58–68. <http://dx.doi.org/10.1016/j.marpolbul.2009.09.001>.
- [137] Dauvin JC. Benthic time-series observations in North-eastern European Marine Station: is a European label needed?. *Mar Pollut Bull* 2010;60:483–8. <http://dx.doi.org/10.1016/j.marpolbul.2010.03.002>.
- [138] Frid CLJ, Garwood PR, Robinson L a. Observing change in a North Sea benthic system: a 33 year time series. *J Mar Syst* 2009;77:227–36. <http://dx.doi.org/10.1016/j.jmarsys.2008.01.011>.