

## UHI Research Database pdf download summary

### Scotland's forgotten carbon:

Smeaton, Craig; Austin, William EN; Davies, Althea; Howe, John; Baltzer, Agnes

*Published in:*  
Biogeosciences

*Publication date:*  
2017

*Publisher rights:*  
© Author(s) 2017. CC Attribution 3.0 License.

*The re-use license for this item is:*  
CC BY

*The Document Version you have downloaded here is:*  
Publisher's PDF, also known as Version of record

*The final published version is available direct from the publisher website at:*  
[10.5194/bg-14-5663-2017](https://doi.org/10.5194/bg-14-5663-2017)

### [Link to author version on UHI Research Database](#)

*Citation for published version (APA):*

Smeaton, C., Austin, W. EN., Davies, A., Howe, J., & Baltzer, A. (2017). Scotland's forgotten carbon: a national assessment of mid-latitude fjord sedimentary carbon stocks. *Biogeosciences*, 14(24), 5663-5674.  
<https://doi.org/10.5194/bg-14-5663-2017>

#### General rights

Copyright and moral rights for the publications made accessible in the UHI Research Database are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights:

- 1) Users may download and print one copy of any publication from the UHI Research Database for the purpose of private study or research.
- 2) You may not further distribute the material or use it for any profit-making activity or commercial gain
- 3) You may freely distribute the URL identifying the publication in the UHI Research Database

#### Take down policy

If you believe that this document breaches copyright please contact us at [RO@uhi.ac.uk](mailto:RO@uhi.ac.uk) providing details; we will remove access to the work immediately and investigate your claim.



## **Sedimentary Carbon Stocks: A National Assessment of Scotland's Fjords.**

Craig Smeaton (1), William Austin (1,2), Althea Davies (1), and John Howe (2)

(1) University of St-Andrews, Department of Geography & Sustainable Development, St-Andrews, United Kingdom (cs244@st-andrews.ac.uk), (2) Scottish Association for Marine Science, Scottish Marine Institute, Dunbeg, Oban, PA37 1QA

Coastal sediments have been shown to be globally significant repositories for carbon (C) with an estimated 126.2 Tg of C being buried annually (Duarte et al. 2005). Though it is clear these areas are important for the long-term storage of C the actual quantity of C held within coastal sediment remains largely unaccounted for. The first step to understanding the role the coastal ocean plays in the global C cycle is to quantify the C held within these coastal sediments.

Of the different coastal environment fjords have been shown to be hotspots for C burial with approximately 11 % of the annual global marine carbon sequestration occurring within fjordic environments (Smith et al. 2015). Through the development of a joint geophysical and geochemical methodology we estimated that the sediment in a mid-latitude fjord holds  $26.9 \pm 0.5$  Mt of C (Smeaton et al., 2016), with these results suggesting that Scottish mid-latitude fjords could be a significant unaccounted store of C equivalent to their terrestrial counterparts (i.e. peatlands).

Through the application of the joint geophysical and geochemical methodology developed by Smeaton et al (2016) to a number of other mid-latitude fjords, we will create detailed estimations of the sedimentary C stored at these individual sites. Using these detailed C stock estimations in conjunction with upscaling techniques we will establish the first national estimation of fjordic sedimentary C stocks. The data produced will allow for the sedimentary C stocks to be compared to other national C stocks, such as the Scottish peatlands (Chapman et al. 2009) and forestry (Forestry Commission, 2016).

Alongside quantifying this large unaccounted for store of C in the coastal ocean this work also lays foundations for future work to understand the role of the coastal ocean in the global C cycle.

Duarte, C. M., Middelburg, J. J., and Caraco, N.: Major role of marine vegetation on the oceanic carbon cycle, *Biogeosciences*, 2, 1–8, doi:10.5194/bg-2-1-2005, 2005.

Smeaton, C., Austin, W. E. N., Davies, A. L., Baltzer, A., Abell, R. E., and Howe, J. A.: Substantial stores of sedimentary carbon held in mid-latitude fjords, *Biogeosciences*, 13, 5771–5787, doi:10.5194/bg-13-5771-2016, 2016.

Smith, R. W., Bianchi, T. S., Allison, M., Savage, C., and Galy, V.: High rates of organic carbon burial in fjord sediments globally, *Nat. Geosci.*, 8, 450–453, doi:10.1038/NGEO2421, 2015.