

## UHI Research Database pdf download summary

### Increased occurrence of the jellyfish *Periphylla periphylla* in the European high Arctic

Geoffroy, Maxime; Berge, Jørgen; Majaneva, Sanna; Johnsen, Geir; Langbehn, Tom J.; Cottier, Finlo; Mogstad, Aksel Alstad; Zolich, Artur; Last, Kim

*Published in:*  
Polar Biology

*Publication date:*  
2018

*The re-use license for this item is:*  
CC BY-NC

*The Document Version you have downloaded here is:*  
Peer reviewed version

*The final published version is available direct from the publisher website at:*  
[10.1007/s00300-018-2368-4](https://doi.org/10.1007/s00300-018-2368-4)

### [Link to author version on UHI Research Database](#)

*Citation for published version (APA):*

Geoffroy, M., Berge, J., Majaneva, S., Johnsen, G., Langbehn, T. J., Cottier, F., Mogstad, A. A., Zolich, A., & Last, K. (2018). Increased occurrence of the jellyfish *Periphylla periphylla* in the European high Arctic. *Polar Biology*, 41(12), 2615-2619. <https://doi.org/10.1007/s00300-018-2368-4>

#### General rights

Copyright and moral rights for the publications made accessible in the UHI Research Database are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights:

- 1) Users may download and print one copy of any publication from the UHI Research Database for the purpose of private study or research.
- 2) You may not further distribute the material or use it for any profit-making activity or commercial gain
- 3) You may freely distribute the URL identifying the publication in the UHI Research Database

#### Take down policy

If you believe that this document breaches copyright please contact us at [RO@uhi.ac.uk](mailto:RO@uhi.ac.uk) providing details; we will remove access to the work immediately and investigate your claim.

1 **Increased occurrence of the jellyfish *Periphylla periphylla* in the European high**  
2 **Arctic**

3

4 Maxime Geoffroy<sup>1,2\*</sup>, Jørgen Berge<sup>2,3,4</sup>, Sanna Majaneva<sup>2,4</sup>, Geir Johnsen<sup>3,4</sup>, Tom J.  
5 Langbehn<sup>5</sup>, Finlo Cottier<sup>6,2</sup>, Aksel Alstad Mogstad<sup>4</sup>, Artur Zolich<sup>7</sup>, Kim Last<sup>6</sup>

6

7 <sup>1</sup>Centre for Fisheries Ecosystems Research, Marine Institute of Memorial University of  
8 Newfoundland, St. John's, Canada

9 <sup>2</sup>UiT The Arctic University of Norway, Department of Arctic and Marine Biology,  
10 Tromsø, Norway

11 <sup>3</sup>University Centre in Svalbard, Longyearbyen, Norway

12 <sup>4</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of  
13 Biology, Norwegian University of Science and Technology, Trondheim, Norway

14 <sup>5</sup>University of Bergen, Department of Biological Sciences, Bergen, Norway

15 <sup>6</sup>Scottish Association for Marine Science, Oban, United Kingdom

16 <sup>7</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of  
17 Engineering Cybernetics, Norwegian University of Science and Technology,  
18 Trondheim, Norway

19

20 \*Corresponding author ([maxime.geoffroy@mi.mun.ca](mailto:maxime.geoffroy@mi.mun.ca); +1-709-778-0499)

21

22 **Abstract**

23 The jellyfish *Periphylla periphylla*, which can have strong ecological impacts on its  
24 environment, is ubiquitous in the Norwegian Sea and its range was predicted to extend  
25 northwards. The abundance of *P. periphylla* in the northern Barents Sea increased since  
26 2014 and, for the first time, several individuals were collected within a high Arctic fjord  
27 ( $>78^{\circ}\text{N}$ ) in western Spitsbergen in January 2017. The low solar irradiance prevailing  
28 during the polar night and an increased inflow of relatively warm Atlantic water in the  
29 European Arctic since the last decade likely provide suitable conditions for the medusa  
30 to colonise Svalbard's fjords during the winter months. However, light avoidance  
31 constrains the photophobic *P. periphylla* to deeper offshore areas during the midnight  
32 sun period. The current occurrence of *P. periphylla* in high Arctic fjords during the  
33 polar night will have a limited impact on marine ecosystems in the short-term, but long-  
34 term effects are more uncertain if its abundance continues to increase.

35

36 **Keywords:** Helmet jellyfish, polar night, Arctic Ocean, Barents Sea, Svalbard,  
37 borealization

38

## 39 **Introduction**

40 The jellyfish *Periphylla periphylla* is a cosmopolitan deep-water species, generally  
41 occurring in low densities ( $<0.02$  individuals  $m^{-3}$ ) at meso- and bathypelagic depths  
42 (Lucas and Reed 2010 and references therein). Yet, under appropriate environmental  
43 conditions *P. periphylla* can also thrive in coastal fjords. For instance, it can reach  
44 exceptionally high densities of  $>2$  individuals  $m^{-3}$  in some Norwegian fjords since the  
45 1970s (e.g. Fosså 1992; Sørnes et al. 2008). High light absorption, resulting in dim light  
46 conditions, and limited water exchanges within the fjords are the main factors behind  
47 blooming populations of this tactile and photophobic predator (Sørnes et al. 2007;  
48 Aksnes et al. 2009; Dupont and Aksnes 2010).

49

50 *Periphylla periphylla* inhabits the Norwegian, Iceland and Greenland Seas (Dalpadado  
51 et al. 1998), but is rarely reported in the northern Barents Sea or in high Arctic fjords  
52 (Gulliksen and Svensen 2004; Gjørseter et al. 2017). Exhaustive reviews of marine  
53 organisms from the Barents Sea and Svalbard archipelago during the last decades  
54 documented the presence of other scyphomedusae, mainly *Cyanea capillata*, but not  
55 that of *P. periphylla* (Zelickman et al. 1972; Palerud et al. 2004). More recently, Tiller  
56 et al. (2017) predicted a northward expansion of the coastal distribution of *P. periphylla*  
57 as a result of climate change (Tiller et al. 2017). In that case, the low solar irradiance  
58 prevailing during the polar night could provide ideal conditions for the jellyfish to  
59 colonise high Arctic fjords in winter.

60

61 Here, we report records of *P. periphylla* in the Barents Sea and in high Arctic fjords of  
62 the Svalbard archipelago, which indicate increased occurrence of the jellyfish in the  
63 European high Arctic in recent years. We further discuss the potential causes and

64 ecological implications of recent occurrence of *P. periphylla* in Svalbard fjords.

65

## 66 **Methods**

### 67 **Biological sampling**

68 We conducted biological sampling in Svalbard in January, May and August 2017, with  
69 a Harstad pelagic trawl (9 × 20-m opening and 1-cm cod end mesh) towed at a given  
70 sampling depth for 20 min at 3 knots by the R/V *Helmer Hanssen* (Online Resource 1).  
71 Samples were sorted and counted on board. *Periphylla periphylla* drifting at the surface  
72 of Kongsfjorden in January were also collected with a dip net (<1-cm mesh) from the  
73 pier in Ny-Ålesund (Fig. 1a). Three of these individuals in good conditions were  
74 dissected for stomach content analyses.

75

76 In addition to the original sampling in 2017, we reviewed annual reports from the  
77 Norwegian Institute of Marine Research (IMR, [www.imr.no](http://www.imr.no)) documenting the  
78 occurrence of marine fish and zooplankton in offshore areas of the Barents Sea from  
79 2005 to 2016. The Institute conducted an average of 291 trawling stations year<sup>-1</sup>  
80 (SD=65) during their fall surveys (August-October; details in Prozorkevich and  
81 Sunnanå 2017). The location of the stations remained similar from one year to another  
82 and covered the entire Barents Sea (Johannesen et al. 2017).

83

84 We also reviewed trawl and plankton datasets from marine biology field courses offered  
85 at the University Centre in Svalbard (UNIS) to document the occurrence of *P.*  
86 *periphylla* in the Svalbard area since 2003 (Jørgen Berge, Paul Renaud and Janne  
87 Søreide, UNIS; unpublished data). A minimum of 10 trawl stations year<sup>-1</sup> were sampled  
88 by UNIS, either during fall (2003-2014) or during spring and fall (2015-2016). The

89 stations were concentrated on the western and northern coasts of Svalbard, from  
90 Isfjorden to Rijpfjorden and to the ice-edge of the central Arctic. Both IMR and UNIS  
91 surveys were conducted with a Harstad or Åkra pelagic trawl and a Campbell 1800  
92 bottom trawl, as well as with plankton nets (WP2, WP3 and MIK-net). It is worth noting  
93 that, in the past decades, gelatinous zooplankton were often disregarded during marine  
94 surveys in the Arctic (Raskoff et al. 2005), but IMR and UNIS documented the  
95 occurrence of gelatinous zooplankton. See Prozorkevich and Sunnanå (2017) and  
96 Renaud et al. (2012) for details about IMR and UNIS sampling, respectively.

97

### 98 **Environmental sampling**

99 We used moored instrumentation to measure water temperature in Kongsfjorden, a  
100 fjord with a maximum depth of 380 m, for the period 2002 until 2017 (Cottier et al.  
101 2005; Berge et al. 2015). The mooring was located in water depths of 200-250 m in the  
102 outer part of Kongsfjorden, which provided a direct connection with the shelf waters  
103 (Cottier et al. 2007). It included 10 temperature sensors (manufactured by either Seabird  
104 or Vemco) positioned from ~20 m below the surface to within 15 m of the seabed. The  
105 precision of the temperature sensors was  $>0.1^{\circ}\text{C}$  after calibration. We calculated the  
106 relationship between years and temperatures and tested for autocorrelation using the R  
107 package nlme, and the best model was selected based on the Akaike information  
108 criterion corrected for small sample size (AICc; Hurvich and Tsai 1993) calculated with  
109 the R package MuMIn.

110

111 A custom-made light sensor deployed at  $77.00^{\circ}\text{N}$  and  $16.35^{\circ}\text{E}$  on January 14, 2018  
112 measured ambient irradiance at 1 m depth (50 m from the research vessel with external  
113 lights off) in the 400–700 nm wavelength range. Measurements were conducted during

114 daytime (15h30 – 16h30 UTC). To obtain absolute irradiance, we calibrated the light  
115 sensor by comparing raw data from the light sensor (digital counts) with absolute values  
116 from a QEPro spectrometer attached to a 2-m long optical fiber (model QP1000-2-vis-  
117 bx) and equipped with a 2pi light collector (model CC-3-UV-S; all from Ocean Optics,  
118 USA) when exposed to a projector at different light intensities in a darkroom (range  
119  $0.24\text{--}42.8 \times 10^{-5} \text{ W m}^{-2}$ ). The QEPro spectrometer was calibrated for absolute irradiance  
120 using an HL-3-cal calibration lamp connected to a cosine corrector through an optical  
121 fiber (Ocean Optics).

122

## 123 **Results**

### 124 **Occurrence of *Periphylla periphylla* in Svalbard fjords in 2017**

125 In January 2017, *P. periphylla* was present in at least two Svalbard fjords (Fig. 1a). In  
126 Kongsfjorden, four were sampled at 336 – 349 m depth and eight were observed drifting  
127 at the surface (Online Resource 1). Five of the latter, all with a bell diameter  $>6.5$  cm,  
128 were collected from the pier in Ny-Ålesund (Fig. 1a). In Rijpfjorden, one *P. periphylla*  
129 was caught at 223 m. In May 2017, a single specimen was captured at 236 m depth in  
130 Isfjorden (Fig. 1b). No *P. periphylla* were caught in August 2017 (Fig. 1c). The  
131 dissected specimens captured at the surface had a low number of prey in their stomach  
132 (average of 5 prey individual<sup>-1</sup>) and a varied diet consisting of copepods (27% of prey  
133 abundance), pteropods (23%), amphipods (20%), euphausiids (17%), and chaetognaths  
134 (13%).

135

### 136 **Review of existing datasets**

137 The IMR collected scyphomedusae, identified as *Cyanea capillata* and *Aurelia aurita*,  
138 in 2006 and 2008-2013. Yet, no *P. periphylla* were reported from 2005 to 2013. In

139 2014, 29 specimens of *P. periphylla* were captured and, since then, the jellyfish has  
140 been caught annually during IMR's surveys (Fig. 1d and Online Resource 2). No *P.*  
141 *periphylla* were collected by UNIS between 2003 and 2015, but one was caught in 2016  
142 (Online Resource 2).

143

#### 144 **Environmental conditions in western Svalbard**

145 Mean water temperature during polar night (November-February) in Kongsfjorden  
146 increased significantly from 0.3°C (2004) to 4°C (2017) (Fig. 2). The significance of  
147 the regression did not vary when tested for autocorrelation (i.e. p-value remained 0.001)  
148 and the linear non-correlated regression model provided the best fit of the data (AICc  
149 of 37.2 vs. 47.1 for the autocorrelated model). Mean fall (August-October)  
150 temperatures in 2017 were the highest recorded during any year, but did not  
151 significantly increase between 2002 and 2017 despite a variation between 1.7°C and  
152 4.6°C (p=0.14; Fig. 2). Ambient irradiance at 77°N during the polar night (January)  
153 remained  $<3.6 \times 10^{-6} \text{ W m}^{-2}$  ( $<1.66 \times 10^{-5} \text{ } \mu\text{mol quanta m}^{-2} \text{ s}^{-1}$ ) at 1 m depth.

154

#### 155 **Discussion**

156 The occurrence of *Periphylla periphylla* increased in the European high Arctic since  
157 2014. Apart from a non-georeferenced mention of *Periphylla periphylla* in Svalbard  
158 (Gulliksen and Svensen 2004) based on occasional encounters during scientific diving  
159 expeditions (Bjørn Gulliksen; personal communication), the jellyfish was not reported  
160 in the northern Barents Sea prior to 2014. Fosså (1992) documented occasional  
161 occurrence in the southern Barents Sea ( $<76.5^\circ\text{N}$ ), but not in Svalbard. From 2014 to  
162 2016, however, it was annually sampled offshore, west and north of Svalbard (Fig. 1d).  
163 In high Arctic fjords, a first specimen was collected in Kongsfjorden in January 2016



164 (Online Resource 2), and we observed high numbers at the same location for the first  
165 time in January 2017. The West Spitsbergen Current (WSC) can transport organisms  
166 from the Norwegian Sea to the European high Arctic in <1 year (Berge et al. 2005;  
167 Gjørseter et al. 2017). Similarly to other boreal species, it is highly probable that *P.*  
168 *periphylla*, a species ubiquitous to the Greenland and Norwegian seas (Dalpadado et al.  
169 1998), was advected with the more persistent inflow of Atlantic water entering western  
170 Svalbard fjords since 2006 (e.g. Willis et al. 2008).

171

172 During the polar night, we measured that irradiance in western Svalbard fjords remains  
173  $<1.66 \times 10^{-5}$   $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$  at 1 m depth, which is exactly within the range of light  
174 preferences of *P. periphylla* (i.e.  $10^{-7}$ – $5 \times 10^{-3}$   $\mu\text{mol quanta m}^{-2} \text{s}^{-1}$ ; Bozman et al. 2017).  
175 Despite northwards advection within the WSC and favorable ambient irradiance  
176 conditions, most *P. periphylla* avoided western Svalbard fjords until winter 2017. Until  
177 then, winter temperatures remained below their known temperature tolerance range of  
178 4 to 19.8°C (Arai 1997 and references therein). Although low temperature has never  
179 been proven to limit the occurrence of *P. periphylla* and the species inhabits cold waters  
180 near 0°C in Antarctica (Larson et al. 1986), in the northern hemisphere, high  
181 abundances occur in fjords where water temperature remains  $>4^\circ\text{C}$  (e.g. Jarms et al.  
182 2002; Sørnes et al. 2007, 2008; Bozman et al. 2017). We thus suggest that the unique  
183 combination of low irradiance and higher temperature which prevailed in January 2017  
184 (for the first time, mean water temperature reached 4°C during the polar night; Fig. 2)  
185 allowed *P. periphylla* to temporarily colonise Kongsfjorden.

186

187 The phototoxic porphyrin pigments of *P. periphylla* cause them potentially lethal  
188 lesions when they are exposed to light (Jarms et al. 2002), and they generally avoid

189 surface waters with higher irradiance during daytime (Kaartvedt et al. 2007; Bozman  
190 et al. 2017). Hence, surface aggregations of *P. periphylla* are only possible during the  
191 polar night in the high Arctic. During the midnight sun period, *P. periphylla* needs to  
192 descend to depth and likely prefers deeper aphotic offshore regions to coastal areas,  
193 which would explain why only one specimen was collected in May and none in August  
194 (Fig. 1b,c). Scyphozoan jellyfish conduct vertical migrations, but also exhibit  
195 horizontal active swimming behavior (Moriarty et al. 2012; Kaartvedt et al. 2015).  
196 *Periphylla periphylla* could thus descend to depth and/or actively avoid Kongsfjorden  
197 from March onwards, prior to the midnight sun period.

198

#### 199 **Potential ecological impacts of *P. periphylla* occurrence in high Arctic fjords**

200 Despite an increasing number of *P. periphylla* in the European high Arctic, its current  
201 abundance remains low compared to blooming populations further south. Acoustic  
202 surveys conducted in Kongsfjorden in January 2017 suggested that the abundance of *P.*  
203 *periphylla* in the top 100 m was  $<0.07$  individuals  $m^{-2}$  (Online Resource 3). Moreover,  
204 *P. periphylla* has a varied diet and a low predation rate (i.e. 1-34 prey  $day^{-1}$ ; Youngbluth  
205 and Båmstedt 2001). Hence, the occurrence of *P. periphylla* in high Arctic fjords during  
206 polar night will likely have a limited impact on marine ecosystems in the short-term. If  
207 low temperature limits the presence of *P. periphylla* in the high Arctic, the ongoing  
208 increase in water temperatures could result in higher abundance, and higher impact, in  
209 the long-term.

210

211 **Acknowledgements.** Pedro de la Torre and Daniel Vogedes provided logistic supports.  
212 Elena Eriksen provided valuable information regarding IMR samples. Bodil Bluhm,  
213 Marine Cusa, Néstor Santana Hernandez, Janne Søreide, and Malin Daase contributed

214 to sampling in 2017. Calum Bachell, Dora Marie Alvsvåg, Øystein Sture, Kertu Liis  
215 Krigul, Jonathan Björklund contributed to fieldwork in January 2017 through the  
216 University Center in Svalbard course AB334/834. A particular thanks to Colin Griffiths  
217 for overseeing the collection of mooring data in Kongsfjorden since 2002.

218

219 **Funding.** The study was co-funded by two projects supported by the Research Council  
220 of Norway (NFR): Arctic ABC (#244319) and AMOS (#223254), and by the  
221 Norwegian Biodiversity Information Centre project GooseAlien (#47-15-70184235).  
222 The mooring work was supported by the UK Natural Environment Research Council  
223 Oceans 2025 and Northern Sea Program with further support from NFR projects Circa  
224 (#214271), Cleopatra (#178766), Cleopatra II (#216537), and Marine Night (#226471).  
225 TJJ acknowledges funding through the MARmaED project, that received funding from  
226 the European Union's Horizon 2020 research and innovation programme under the  
227 Marie Skłodowska-Curie grant agreement No 675997. This study is a contribution  
228 towards the ARCTOS research network ([www.arctos.uit.no](http://www.arctos.uit.no)), the Arctic Science  
229 Partnership ([ww.asp.org](http://ww.asp.org)) and the Ocean Frontier Institute funded through Canada's  
230 First Research Excellence Fund. The results of this publication reflect only the authors'  
231 view and funding agencies are not responsible for any use that may be made of the  
232 information it contains.

233

234 **Compliance with ethical standards**

235 **Conflict of interests.** The authors declare that they have no conflict of interests.

236

237 **Ethical approval.** All applicable international, national, and/or institutional guidelines  
238 for the care and use of animals were followed. This study was carried out with

239 permission from the Governor on Svalbard and followed the strict regulations regarding  
240 health, environment and safety enforced at UNIS and UiT.

241

## 242 **References**

243 Aksnes DL, Dupont N, Staby A, Fiksen Ø, Kaartvedt S, Aure J (2009) Coastal water

244 darkening and implications for mesopelagic regime shifts in Norwegian fjords.

245 Mar Ecol Prog Ser 387:39–49. <https://doi.org/10.3354/meps08120>

246 Arai MN (1997) A functional biology of Scyphozoa. Chapman & Hall, London

247 Berge J, Johnsen G, Nilsen F, Gulliksen B, Slagstad D (2005) Ocean temperature

248 oscillations enable reappearance of blue mussels *Mytilus edulis* in Svalbard after

249 a 1000 year absence. Mar Ecol Prog Ser 303:167–175

250 Berge J, Hegglund K, Lønne OJ et al (2015) First records of Atlantic Mackerel

251 (*Scomber scombrus*) from the Svalbard Archipelago, Norway, with possible

252 explanations for the extension of its distribution. Arctic 68:54–61.

253 <https://dx.doi.org/10.14430/arctic4455>

254 Bozman A, Titelman J, Kaartvedt S, Eiane K, Aksnes DL (2017) Jellyfish distribute

255 vertically according to irradiance. J Plankton Res 39:280–289. [https://doi.org/](https://doi.org/10.1093/plankt/fbw097)

256 [10.1093/plankt/fbw097](https://doi.org/10.1093/plankt/fbw097)

257 Cottier FR, Nilsen F, Inall ME, Gerland S, Tverberg V, Svendsen H (2007)

258 Wintertime warming of an Arctic shelf in response to large-scale atmospheric

259 circulation. Geophys Res Lett 34:L10607. [https://doi.org/10.1029/](https://doi.org/10.1029/2007GL029948)

260 [2007GL029948](https://doi.org/10.1029/2007GL029948)

261 Cottier F, Tverberg V, Inall M, Svendsen H, Nilsen F, Griffiths C (2005) Water mass

262 modification in an Arctic fjord through cross-shelf exchange: The seasonal

263 hydrography of Kongsfjorden, Svalbard. J Geophys Res 110:C12005. [https://](https://doi.org/10.1029/2004JC002757)  
264 [doi:10.1029/2004JC002757](https://doi.org/10.1029/2004JC002757)

265 Dalpadado P, Ellertsen B, Melle W, Skjoldal HR (1998) Summer distribution patterns  
266 and biomass estimates of macrozooplankton and micronekton in the Nordic  
267 Seas. Sarsia 83:103–116. <https://doi.org/10.1080/00364827.1998.10413676>

268 Dupont N, Aksnes DL (2010) Simulation of optically conditioned retention and mass  
269 occurrences of *Periphylla periphylla*. J Plankton Res 32:773–783. [https://](https://doi.org/10.1093/plankt/fbq015)  
270 [doi.org/10.1093/plankt/fbq015](https://doi.org/10.1093/plankt/fbq015)

271 Fosså JH (1992) Mass occurrence of *Periphylla periphylla* (Scyphozoa, Coronatae) in  
272 a Norwegian fjord. Sarsia 77:237–251. [https://doi.org/10.1080/](https://doi.org/10.1080/00364827.1992.10413509A)  
273 [00364827.1992.10413509A](https://doi.org/10.1080/00364827.1992.10413509A)

274 Gjørseter H, Wiebe PH, Knutsen T, Ingvaldsen RB (2017) Evidence of diel vertical  
275 migration of mesopelagic sound-scattering organisms in the Arctic. Front Mar  
276 Sci 4:1–14. <https://doi.org/10.3389/fmars.2017.00332>

277 Gulliksen B, Svensen E (2004) Svalbard and life in polar oceans. Kom forlag,  
278 Kristiansund

279 Hurvich CM, Tsai CL (1993) A corrected Akaike information criterion for vector  
280 autoregressive model selection. J time series anal 14: 271–279.  
281 <https://doi.org/10.1111/j.1467-9892.1993.tb00144.x>

282 Jarms G, Tiemann H, Båmstedt U (2002) Development and biology of *Periphylla*  
283 *periphylla* (Scyphozoa: Coronatae) in a Norwegian fjord. Mar Biol 141:647–  
284 657. <https://doi.org/10.1007/s00227-002-0858-x>

285 Johannesen E, Mørk HL, Korsbrekke K et al (2017) Arctic fishes in the Barents Sea  
286 2004-2015: Changes in abundance and distribution, IMR/PINRO Joint Report  
287 Series, No. 1/2017

288 Kaartvedt S, Klevjer TA, Torgersen T, Sørnes TA, Røstad A (2007) Diel vertical  
289 migration of individual jellyfish (*Periphylla periphylla*). *Limn Oceanogr*  
290 52:975–983. <https://doi.org/10.4319/lo.2007.52.3.0975>

291 Kaartvedt S, Ugland KI, Klevjer TA, Røstad A, Titelman J, Solberg I (2015) Social  
292 behaviour in mesopelagic jellyfish. *Sci Rep* 5:1–8. <https://doi.org/10.1038/srep11310>

294 Larson RJ (1986) Pelagic Scyphomedusae (Scyphozoa: Coronatae and  
295 Semaestomae) of the Southern Ocean. *Biology of the Antarctic Seas XVI*.  
296 *Antarct Res Ser* 41:59-165

297 Lucas CH, Reed AJ (2010) Gonad morphology and gametogenesis in the deep-sea  
298 jellyfish *Atolla wyvillei* and *Periphylla periphylla* (Scyphozoa: Coronatae)  
299 collected from Cape Hatteras and the Gulf of Mexico. *J Mar Biol Ass UK* 90:  
300 1095–1104. <https://doi.org/10.1017/S0025315409000824>

301 Moriarty P, Andrews K, Harvey C, Kawase M (2012) Vertical and horizontal  
302 movement patterns of scyphozoan jellyfish in a fjord-like estuary. *Mar Ecol*  
303 *Prog Ser* 455:1–12. <https://doi.org/10.3354/meps09783>

304 Pagès F, Kurbjewit F (1994) Vertical distribution and abundance of mesoplanktonic  
305 medusae and siphonophores from the Weddell Sea, Antarctica. *Polar Biol*  
306 14:243–251. <https://doi.org/10.1007/BF00239172>

307 Palerud R, Gulliksen B, Brattegard T, Sneli JA, Vader W (2004) The marine macro-  
308 organisms in Svalbard waters. In: Presterud P, Strøm H, Goldman HV (eds) *A*  
309 *catalogue of the terrestrial and marine animals of Svalbard*, vol 201, 1st edn.  
310 Norsk Polarinstitutt Skrifter, Tromsø, pp 5–56

311 Prozorkevich D, Sunnanå K (2017) Survey report from the joint Norwegian/Russian  
312 ecosystem survey in the Barents Sea and adjacent waters, August-October 2016,  
313 IMR/PINRO Joint Report Series, No. 1/2017

314 Raskoff KA, Purcell JE, Hopcroft RR (2005) Gelatinous zooplankton of the Arctic  
315 Ocean: *in situ* observations under the ice. *Polar Biol* 28:207-217. <https://doi.org/10.1007/s00300-004-0677-2>

316

317 Renaud PE, Berge J, Varpe Ø, Lønne OJ, Nahrgang J, Ottesen C, Hallanger I (2012)  
318 Is the poleward expansion by Atlantic cod and haddock threatening native polar  
319 cod, *Boreogadus saida*? *Polar Biol* 35:401–412. [https://doi.org/10.1007/s00300-](https://doi.org/10.1007/s00300-011-1085-z)  
320 [011-1085-z](https://doi.org/10.1007/s00300-011-1085-z)

321 Sørnes TA, Aksnes DL, Båmstedt U, Youngbluth MJ (2007) Causes for mass  
322 occurrences of the jellyfish *Periphylla periphylla*: A hypothesis that involves  
323 optically conditioned retention. *J Plankton Res* 29:157–167. [https://doi.org/](https://doi.org/10.1093/plankt/fbm003)  
324 [10.1093/plankt/fbm003](https://doi.org/10.1093/plankt/fbm003)

325 Sørnes TA, Hosia A, Båmstedt U, Aksnes DL (2008) Swimming and feeding in  
326 *Periphylla periphylla* (Scyphozoa, Coronatae). *Mar Biol* 153:653–659.  
327 <https://doi.org/10.1007/s00227-007-0839-1>

328 Tiller RG, Borgersen ÅL, Knutsen Ø et al (2017) Coming soon to a fjord near you:  
329 Future jellyfish scenarios in a changing climate. *Coast Manag* 45:1–23.  
330 <https://doi.org/10.1080/08920753.2017.1237239>

331 Willis KJ, Cottier FR, Kwaśniewski S (2008) Impact of warm water advection on the  
332 winter zooplankton community in an Arctic fjord. *Polar Biol* 31:475–481.  
333 <https://doi.org/10.1007/s00300-007-0373-0>

334 Youngbluth MJ, Båmstedt U (2001) Distribution, abundance, behavior and  
335 metabolism of *Periphylla periphylla*, a mesopelagic coronate medusa in a

336 Norwegian fjord. *Hydrobiologia* 451:321–333. <https://doi.org/10.1023/>

337 A:1011874828960


338 Zelickman EA (1972) Distribution and ecology of the pelagic hydromedusae,

339 siphonophores and ctenophores of the Barents Sea, based on perennial plankton

340 collections. *Mar Biol* 17:256–264. <https://doi.org/10.1007/BF00366301>

341

342 **Figure captions**



343


344 **Fig. 1** (a-c) General map (insert) and survey area in Svalbard. Stations without (empty

345 circle) and with (black dot) *P. periphylla* are indicated. The white star indicates surface

346 observations in Ny-Ålesund. (d) Previous observations by year. The numbers of *P.*

347 *periphylla* sampled at each location are included

348



349

350 **Fig. 2** Mean fjord temperatures measured from 20 m to bottom in Kongsfjorden during

351 fall (August-October; triangles) and the polar night (November-February; dots)



**Increased occurrence of the jellyfish *Periphylla periphylla* in the European high Arctic**

Maxime Geoffroy<sup>1,2\*</sup>, Jørgen Berge<sup>2,3,4</sup>, Sanna Majaneva<sup>2,4</sup>, Geir Johnsen<sup>3,4</sup>, Tom J. Langbehn<sup>5</sup>, Finlo Cottier<sup>6,2</sup>, Aksel Alstad Mogstad<sup>4</sup>, Artur Zolich<sup>7</sup>, Kim Last<sup>6</sup>

<sup>1</sup>Centre for Fisheries Ecosystems Research, Marine Institute of Memorial University of Newfoundland, St. John's, Canada

<sup>2</sup>UiT The Arctic University of Norway, Department of Arctic and Marine Biology, Tromsø, Norway

<sup>3</sup>University Centre in Svalbard, Longyearbyen, Norway

<sup>4</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of Biology, Norwegian University of Science and Technology, Trondheim, Norway

<sup>5</sup>University of Bergen, Department of Biological Sciences, Bergen, Norway

<sup>6</sup>Scottish Association for Marine Science, Oban, United Kingdom

<sup>7</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of Engineering Cybernetics, Norwegian University of Science and Technology, Trondheim, Norway

\*Corresponding author ([maxime.geoffroy@mi.mun.ca](mailto:maxime.geoffroy@mi.mun.ca); +1-709-778-0499)

**Electronic Supplementary Material 1**

**Supplementary Table ESM1.** Details of sampling conducted in Svalbard in January, May and August 2017.

Date	Latitude (°N)	Longitude (°E)	Sampling Depth (m)	Type	# of <i>P. periphylla</i>
2017-01-09	79.016	11.177	322	Pelagic trawl	0
2017-01-10	79.085	11.434	336	Pelagic trawl	3
2017-01-10	79.091	11.442	349	Pelagic trawl	1
2017-01-11	80.390	13.533	206	Pelagic trawl	0
2017-01-11	80.378	12.472	1041	Pelagic trawl	0
2017-01-13	81.218	14.364	2194	Pelagic trawl	0
2017-01-14	80.570	17.397	354	Pelagic trawl	0
2017-01-14	80.202	22.117	223	Pelagic trawl	1
2017-01-16	79.444	11.061	211	Pelagic trawl	0
2017-01-19	78.936	11.915	0	Dip net	1
2017-01-21	78.936	11.915	0	Dip net	4
2017-01-21	78.936	11.915	0	Visual obs.	3
2017-05-08	78.177	15.063	236	Pelagic trawl	1
2017-05-09	78.583	11.551	266	Pelagic trawl	0
2017-05-11	79.502	10.046	401	Pelagic trawl	0
2017-05-12	79.387	9.338	302	Pelagic trawl	0
2017-08-08	79.003	11.370	215	Pelagic trawl	0
2017-08-08	79.006	11.279	380	Pelagic trawl	0
2017-08-09	78.577	11.567	335	Pelagic trawl	0
2017-08-11	79.480	12.010	215	Pelagic trawl	0
2017-08-12	80.192	22.140	230	Pelagic trawl	0
2017-08-13	80.067	22.109	164	Pelagic trawl	0
2017-08-17	78.231	15.281	226	Pelagic trawl	0

**Increased occurrence of the jellyfish *Periphylla periphylla* in the European high Arctic**

Maxime Geoffroy<sup>1,2\*</sup>, Jørgen Berge<sup>2,3,4</sup>, Sanna Majaneva<sup>2,4</sup>, Geir Johnsen<sup>3,4</sup>, Tom J. Langbehn<sup>5</sup>, Finlo Cottier<sup>6,2</sup>, Aksel Alstad Mogstad<sup>4</sup>, Artur Zolich<sup>7</sup>, Kim Last<sup>6</sup>

<sup>1</sup>Centre for Fisheries Ecosystems Research, Marine Institute of Memorial University of Newfoundland, St. John's, Canada

<sup>2</sup>UiT The Arctic University of Norway, Department of Arctic and Marine Biology, Tromsø, Norway

<sup>3</sup>University Centre in Svalbard, Longyearbyen, Norway

<sup>4</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of Biology, Norwegian University of Science and Technology, Trondheim, Norway

<sup>5</sup>University of Bergen, Department of Biological Sciences, Bergen, Norway

<sup>6</sup>Scottish Association for Marine Science, Oban, United Kingdom

<sup>7</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of Engineering Cybernetics, Norwegian University of Science and Technology, Trondheim, Norway

\*Corresponding author ([maxime.geoffroy@mi.mun.ca](mailto:maxime.geoffroy@mi.mun.ca); +1-709-778-0499)

**Electronic Supplementary Material 2**

**Table ESM2.** Details of *P. periphylla* previously sampled in the European high Arctic. NA indicates information that is not available.

Date	Latitude (°N)	Longitude (°E)	Area	Sampling device	Sampling depth (m)	Sampling duration (min.)	# of <i>P. periphylla</i>	Reference
2017-04-12	78.250	15.45	Adventfjorden	Visual obs.	0	-	1	a
2016-09	71.783	21.783	South of Svalbard	Pelagic trawl	NA	NA	1	b
2016-09	73.000	18.417	South of Svalbard	Pelagic trawl	NA	NA	1	b
2016-09	75.400	14.367	South of Svalbard	Pelagic trawl	NA	NA	1	b
2016-08-28	80.828	15.880	North slope	Pelagic trawl	550	NA	1	c
2016-02	78.936	11.915	Kongsfjorden	Visual obs.	0	-	1	d
2015-09-24	72.970	14.680	South of Svalbard	Pelagic trawl	NA	71	1	b
2015-09-16	73.000	28.470	South of Svalbard	Pelagic trawl	NA	31	1	b

2015-09-14	76.450	35.150	West of Svalbard	Bottom trawl	242	15	1	b
2015-09-13	75.320	34.570	South of Svalbard	Pelagic trawl	NA	92	1	b
2015-09-05	78.650	9.070	West of Svalbard	Pelagic trawl	NA	36	1	b
2015-09-04	79.700	9.070	West of Svalbard	Bottom trawl	404	33	1	b
2015-09-03	75.320	28.120	South of Svalbard	Bottom trawl	285	16	1	b
2015-09-03	79.759	6.350	West of Svalbard	Pelagic trawl	350-450	29	0.348 kg nmi <sup>-2</sup>	e
2015-09-02	79.750	6.350	West of Svalbard	Bottom trawl	1014	70	1	b
2015-08-30	80.880	6.870	North slope	Bottom trawl	917	55	1	b
2015-08-29	80.420	8.400	North slope	Bottom trawl	861	45	1	b
2015-08-26	81.080	17.100	North slope	Bottom trawl	806	53	2	b
2015-08-23	76.350	12.700	West of Svalbard	Pelagic trawl	450	68	1	b
2015-08-23	81.561	15.294	North slope	Pelagic trawl	258-400	30	0.492 kg nmi <sup>-2</sup>	e
2015-08-22	80.750	15.520	North slope	Bottom trawl	1632	155	1	b
2015-08-20	80.250	16.780	North slope	Bottom trawl	297	15	1	b
2015-08-18	78.020	9.380	West of Svalbard	Bottom trawl	577	127	1	b
2014-09-12	77.482	11.133	West of Svalbard	Pelagic trawl	NA	30	1	b
2014-09-11	76.898	11.933	West of Svalbard	Pelagic trawl	NA	15	1	b
2014-09-10	75.352	14.071	South of Svalbard	Pelagic trawl	NA	30	1	b
2014-09-10	76.406	12.785	West of Svalbard	Pelagic trawl	NA	30	2	b
2014-09-08	73.477	13.943	South of Svalbard	Pelagic trawl	NA	30	1	b
2014-09-07	71.799	15.518	South of Svalbard	Pelagic trawl	NA	30	1	b
2014-09-03	78.617	6.615	West of Svalbard	Bottom trawl	NA	141	1	b
2014-09-02	78.602	9.123	West of Svalbard	Bottom trawl	NA	65	9	b
2014-09-02	78.616	8.610	West of Svalbard	Bottom trawl	NA	63	1	b
2014-08-23	79.644	5.344	West of Svalbard	Bottom trawl	NA	120	1	b
2014-08-20	71.677	23.167	South of Svalbard	Bottom trawl	NA	16	10	b

References: a) Larissa Beumer, UNIS; Personnal communication; b) Elena Eriksen, IMR; Personnal communication c) Paul Renaud, Akvaplan-niva; Personnal communication; d) Leif Arne Håhjem, Norwegian Polar Institute; Personnal communication; e) Gjøsæter et al. (2017)

**Increased occurrence of the jellyfish *Periphylla periphylla* in the European high Arctic**

Maxime Geoffroy<sup>1,2\*</sup>, Jørgen Berge<sup>2,3,4</sup>, Sanna Majaneva<sup>2,4</sup>, Geir Johnsen<sup>3,4</sup>, Tom J. Langbehn<sup>5</sup>, Finlo Cottier<sup>6,2</sup>, Aksel Alstad Mogstad<sup>4</sup>, Artur Zolich<sup>7</sup>, Kim Last<sup>6</sup>

<sup>1</sup>Centre for Fisheries Ecosystems Research, Marine Institute of Memorial University of Newfoundland, St. John's, Canada

<sup>2</sup>UiT The Arctic University of Norway, Department of Arctic and Marine Biology, Tromsø, Norway

<sup>3</sup>University Centre in Svalbard, Longyearbyen, Norway

<sup>4</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of Biology, Norwegian University of Science and Technology, Trondheim, Norway

<sup>5</sup>University of Bergen, Department of Biological Sciences, Bergen, Norway

<sup>6</sup>Scottish Association for Marine Science, Oban, United Kingdom

<sup>7</sup>Centre for Autonomous Marine Operations and Systems (AMOS), Department of Engineering Cybernetics, Norwegian University of Science and Technology, Trondheim, Norway

\*Corresponding author ([maxime.geoffroy@mi.mun.ca](mailto:maxime.geoffroy@mi.mun.ca); +1-709-778-0499)

**Electronic Supplementary Material 3.** Details of acoustic analyses conducted in Kongsfjorden in January 2017.

**Sampling and analysis**

To estimate the density of *Periphylla periphylla* in Kongsfjorden, acoustic transects were conducted with a Wideband Autonomous Transceiver<sup>®</sup> (WBAT; 70 kHz producing a chirp from 55 to 90 kHz) and a multifrequency (125, 200, 455, 769 kHz) Acoustic Zooplankton and Fish Profiler<sup>®</sup> (AZFP). The AZFP was mounted in two different configurations, either on an autonomous surface vehicle (January 19) or on the side of a small boat (January 21), both cruising at 2 knots. The WBAT was mounted on the side of a small boat on January 19. Both instruments were calibrated by the manufacturer.

Acoustic data were processed using Echoview<sup>®</sup> 8. The Target Strength frequency distribution (TS in dB re 1 m<sup>2</sup>; Fig. ESM3) was calculated from the WBAT data using the single target detection and fish track algorithms (Tables ESM3a,b). The analysis was limited to the top 100 m because a lower signal-to-noise ratio prevented single detections at a farther range. Volume backscattering strength echograms (Sv in dB re 1 m<sup>-1</sup>) of the WBAT and AZFP (125 kHz only) were then divided in 1-min cells before being echo integrated over the top 100 m. To compensate for noise amplification at depth by the time-varied gain, a time-varied threshold of -125 dB at 1 m was added to the WBAT data and Echoview's Background Noise Removal algorithm (DeRobertis and Higginbottom 2007) was applied to the AZFP data. The Nautical Area Backscattering Coefficient (NASC in m<sup>-2</sup> nmi<sup>-2</sup>) within each 1-min cell was divided by the average cross-section ( $4\pi\sigma_{bs}$ , in m<sup>-2</sup>) estimated from the TS analysis, and multiplied by the relative abundance of signals between -64 and -59 dB (9.8%; Fig. ESM3) to estimate the average density of *P. periphylla* in the top 100 m

during each transect. Single beam transducers as that of the AZFP do not allow accurate TS measurements. Hence, the TS frequency distribution from the WBAT data was used to estimate the abundance of *P. periphylla* from both the WBAT and AZFP transects.

### Results and interpretation

Distinct Gaussian distributions of acoustic TS are known to indicate the occurrence of distinct species and/or size classes (Simmons and MacLennan 2005). Here, a distinct TS distribution between -64 and -59 dB (mean of -61.3 dB) was detected with the WBAT and represented 9.8% of the targets tracked in the top 100 m of Kongsfjorden on January 19 (Fig. ESM3). This TS distribution was assumed to mainly originate from *P. periphylla* individuals similar to those observed at the surface because a mean TS of -60 dB (SD 3.6 dB; n= 598) (Kaartved et al. 2007) and a peak TS of -62 dB (Klevjer et al. 2009) were previously measured at 38 kHz for *P. periphylla* detected at similar distances. Applying the 9.8% ratio to NASC values, we estimated that mean densities of *P. periphylla* could have reached 0.07 ind. m<sup>-2</sup> on January 19, and 0.02 ind. m<sup>-2</sup> on January 21.

Other abundant targets in Kongsfjorden either have lower (zooplankton) or higher (fish) TS. Zooplankton in Kongsfjorden have a TS between -109 and -74 dB (Berge et al. 2014), which corresponds to the distributions to the left of *P. periphylla* targets in the TS frequency histograms (Fig. ESM3). The main pelagic fish species in Kongsfjorden comprise polar cod (*Boreogadus saida*) and capelin (*Mallotus villosus*) (Hop et al. 2002). Recently, juvenile Atlantic redfish (*Sebastes* spp.) have also colonised the fjord. All these fish have a mean TS > -58 dB, corresponding to the distributions to the right of that attributed to *P. periphylla* (Fig. ESM3). Other occasional targets as *Cyanea capillata* or myctophidae could have partly contributed to the acoustic signal attributed to *P. periphylla*. Fish with a pronounced tilt angle could also have contributed to that signal. Hence, our abundance estimates of *P. periphylla* represent maximum values and were likely biased positively.


#### Supplementary Table ESM3a. Parameters used for the single target detection algorithm in Echoview®.

Parameters	Values
Compensated TS threshold	-120 dB
Pulse length determination level	6 dB
Minimum normalize pulse length	0.001
Maximum normalized pulse length	1.5
Beam compensation model	Simrad LOBE
Maximum beam compensation	6 dB
Maximum standard deviation of minor and major axis angles	0.6°

#### Supplementary Table ESM3b. Parameters used for the fish-tracking algorithm in Echoview®.

	Major axis	Minor axis	Range
Alpha	0.7	0.7	0.7
Beta	0.5	0.5	0.5
Exclusion distance (m)	4.0	4.0	0.3
Missed ping expansion %	0	0	100
Weight (%)	30	30	40

Minimum number of single targets and pings in a track were set to 3, and minimum gap between single targets was set to 5 pings. Note that for wideband systems, the value of compensated TS indicates the peak of the TS-frequency curve.



**Supplementary Figure ESM3.** TS frequency histogram presenting the upper 100 m of the WBAT data collected on January 19. The blue line is the kernel density estimates with a bandwidth of 0.5 (i.e. normal distribution approximations). The dashed red lines indicate the limits of the TS frequency distribution attributed to *P. periphylla* (-64 to -59 dB), and the percentage and mean TS of targets within this distribution are indicated.

### References ESM3

- Berge J, Cottier F, Varpe Ø et al (2014) Arctic complexity: A case study on diel vertical migration of zooplankton. *J Plankton Res* 36:1279–1297. <https://doi.org/10.1093/plankt/fbu059>
- De Robertis A, Higginbottom I (2007) A post-processing technique to estimate the signal-to-noise ratio and remove echosounder background noise. *ICES J Mar Sci* 64: 1282–1291. <https://doi.org/10.1093/icesjms/fsm112>
- Hop H, Pearson T, Hegseth EN et al (2002) The marine ecosystem of Kongsfjorden, Svalbard. *Polar Res* 21: 167–208
- Kaartvedt S, Klevjer TA, Torgersen T, Sørnes TA, Røstad A (2007) Diel vertical migration of individual jellyfish (*Periphylla periphylla*). *Limnol Oceanogr* 52:975–983. <https://doi.org/10.4319/lo.2007.52.3.0975>
- Klevjer TA, Kaartvedt S, Båmstedt U (2009) *In situ* behaviour and acoustic properties of the deep living jellyfish *Periphylla periphylla*. *J Plankton Res* 31:793–803. <https://doi.org/10.1093/plankt/fbp036>
- Simmons EJ, MacLennan DN, Simmonds EJ (2005) *Fisheries acoustics*. Chapman & Hall, London