

UHI Research Database pdf download summary

European expansion of the introduced amphipod *Caprella mutica* Schurin 1935

Cook, Elizabeth J.; Jahnke, Marlene; Kerckhof, Francis; Minchin, Dan; Faasse, Marco; Boos, Karin; Ashton, Gail

Published in:
Aquatic Invasions

Publication date:
2007

The re-use license for this item is:
CC BY

The Document Version you have downloaded here is:
Publisher's PDF, also known as Version of record

The final published version is available direct from the publisher website at:
[10.3391/ai.2007.2.4.11](https://doi.org/10.3391/ai.2007.2.4.11)

[Link to author version on UHI Research Database](#)

Citation for published version (APA):

Cook, E. J., Jahnke, M., Kerckhof, F., Minchin, D., Faasse, M., Boos, K., & Ashton, G. (2007). European expansion of the introduced amphipod *Caprella mutica* Schurin 1935. *Aquatic Invasions*, 2(4), 411-421. <https://doi.org/10.3391/ai.2007.2.4.11>

General rights

Copyright and moral rights for the publications made accessible in the UHI Research Database are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights:

- 1) Users may download and print one copy of any publication from the UHI Research Database for the purpose of private study or research.
- 2) You may not further distribute the material or use it for any profit-making activity or commercial gain
- 3) You may freely distribute the URL identifying the publication in the UHI Research Database

Take down policy

If you believe that this document breaches copyright please contact us at RO@uhi.ac.uk providing details; we will remove access to the work immediately and investigate your claim.

European expansion of the introduced amphipod *Caprella mutica* Schurin 1935

Elizabeth J. Cook^{1*}, Marlene Jahnke¹, Francis Kerckhof², Dan Minchin³, Marco Faasse⁴, Karin Boos⁵ and Gail Ashton⁶

¹Scottish Association for Marine Science, Dunstaffnage Marine Laboratory, Oban, Argyll PA37 1QA, UK
E-mail: Elizabeth.Cook@sams.ac.uk

²MUMM, Marine Environmental Management Section, Royal Belgian Institute of Natural Sciences, 3e en 23e Linierregimentsplein, B-8400 Oostende, Belgium, E-mail: francis.kerckhof@mumm.ac.be

³Marine Organism Investigations, 3 Marina Village, Ballina, Killaloe, Co. Clare, Ireland
E-mail: minchin@indigo.ie

⁴National Museum of Natural History, Naturalis, P.O. Box 9517, 2300 RA Leiden, The Netherlands
E-mail: mafaasse@hetnet.nl

⁵Biologische Anstalt Helgoland, Alfred Wegener Institut for Polar- and Marine Research, P.O. Box 180, 27483 Helgoland, Germany,
E-mail: Karin.Boos@awi.de

⁶Smithsonian Environmental Research Centre, 647 Contees Wharf Road, P.O. Box 28, Edgewater MD 21037, USA, E-mail: ashtong@si.edu

*Corresponding author

Received 1 November 2007; accepted in revised form 27 November 2007

Abstract

The amphipod *Caprella mutica* is one of the most rapidly invading species in Europe and has extended its range throughout North Sea and Celtic Sea coasts and the English Channel in less than fourteen years. It was first described from sub-boreal areas of north-east Asia in 1935 and has since spread to both northern and southern hemispheres. The first European record was from The Netherlands in 1994. Since then it has spread within the North Sea and later to the west coast of Scotland and to Ireland. *C. mutica* is frequently associated with man-made structures and is found in abundance on boat hulls, navigation/ offshore buoys, floating pontoons and aquaculture infrastructure. It is highly likely that its dispersal is associated with vessel movements whilst attached to hull fouling. This species is expected to colonise the west coasts of France and Spain and offshore islands in the north-east Atlantic.

Key words: *Caprella mutica*, Europe, caprellid, crustacean, distribution, biological invasion

Introduction

The caprellid amphipod *Caprella mutica* Schurin 1935 (Figure 1) is indigenous to sub-boreal waters of north-east Asia (Peter the Great Bay, Vladivostok, Russia) and was subsequently identified in the neighbouring Possjet Bay, Japan (Vassilenko 1967) and Akkeshi Bay, Japan (Arimoto 1976). The first reports of *C. mutica* outside its native habitat were from the Pacific and Atlantic coasts of North America in the

1970s (Carlton 1979) and 1980s (Marelli 1981, Cohen and Carlton 1995) and a recent review of its global distribution has shown that within 40 years this species has spread throughout the northern hemisphere and it has recently been found for the first time in New Zealand (Ashton et al. 2007a).

C. mutica is one of the largest caprellid amphipods, mature males attain body lengths of up to 50 mm (Nishimura 1995). Females are highly fecund, producing their first brood

approximately 53 days post-hatching at an average body length of 8.5 mm and at a seawater temperature of 13–14°C (Cook et al. 2007). Each female has an average of two sequential broods released at approximately 20 day intervals. The maximum number of recorded hatchlings produced by a single female at a seawater temperature of $13.0 \pm 0.5^\circ\text{C}$ is 82. Juveniles typically emerge from the brood pouch at a body length of approximately 1.3 mm (Cook et al. 2007). *C. mutica* is able to survive for up to 20 days without additional food under laboratory conditions (Cook et al. 2007) and tolerates temperatures of $< 2^\circ\text{C}$ to 28°C and salinities down to 19 psu over a short (48 h) exposure period (Ashton et al. 2007b). *C. mutica* is an aggressive species, out-competing the native European caprellid *Caprella linearis* for space, even at low densities (Shucksmith in press). In the native range, this species is typically associated with either attached or drifting macroalgae, including *Sargassum* spp. or aquaculture structures, such as the ropes used for the culture of the macroalga *Undaria* spp. in Otsuchi Bay (Kawashima et al. 1999). In regions outside its native range, it has been found associated with areas of anthropogenic activity, such as harbours, marinas, navigation buoys and aquaculture sites (Willis et al. 2004, Ashton 2006, Kerckhof et al. 2007).

Carlton (1979) suggested that *C. mutica* arrived on the Pacific and Atlantic coasts of North America either as a result of numerous independent cross-oceanic introductions with oyster spat, or from small scale transport following its first introduction (Carlton 1996). Direct sequencing of mitochondrial DNA indicates that *C. mutica* was introduced to Europe either directly from Asia or from the Atlantic coast of North America (Ashton 2006). Opportunities for the spread of non-native species to Europe across the Atlantic have existed for more than 500 years with early exploration and the subsequent establishment of shipping routes (Stoner et al. 2002). For a little more than a century ships' ballast water has transmitted many different crustacean taxa, including caprellids (Coutts et al. 2003) and in the last half-century ocean barges and oil platforms (or similar structures) have been moved across oceans (Rodríguez and Suárez 2001, Ruiz and Hewitt 2002) and may have introduced *C. mutica* to Europe. The importation of different species of half-grown oysters as deck



Figure 1. Male (top) and female (bottom) *Caprella mutica* (Photo: T. Nickell)

cargo since ~1870s (Loosanoff 1975) and of Pacific oysters (*Crassostrea gigas*) from British Columbia in the 1960s and later directly from Japan in the 1970s into Europe, may also have provided opportunities for introduction (Wolff 2005). *C. mutica* lacks a free-swimming planktonic larval stage and secondary movements are also likely to involve human activities, such as shipping, aquaculture activities and recreational boating or may otherwise be associated with drifting macroalgae (Ashton 2006).

C. mutica was first identified in European waters in 1994 in a harbour at Neeltje Jans in The Netherlands (M. Faasse, pers. obs.). In a review by Ashton et al. (2007a) its general occurrence in Europe is described. In this account, we report all known localities for this species in European waters and discuss the likely dispersal mechanisms involved in its further spread.

Methods and Materials

Published records and unpublished reports were consulted for verifiable material. Researchers with recorded sightings of *C. mutica* were contacted for further information and if possible, specimens were obtained for confirmation of identification using the taxonomic key by Arimoto (1976).

Results

One hundred and twenty-one records of *Caprella mutica* were found for European waters (Annex), including eleven new localities for 2007. Confirmed records for *C. mutica* are from the North Sea, English Channel, Irish Sea, the west

coast of Scotland and Ireland (Figure 2) and range from 49°29'N to 62°22'N and 10°04'W to 8°26'E. While *C. mutica* was not found at all localities surveyed within this range, the region comprising the greatest number of records was the south-western North Sea. There are presently no records for the Baltic Sea, the Iberian Peninsula or the Mediterranean Sea.

The majority of the European records occur in areas with human activity, such as marinas, ports and shipping routes. In Norway and the west coast of Scotland, *C. mutica* is associated with aquaculture activity, principally fish cage netting and mussel lines. This species has also been found on natural substrata within the sub-littoral zone on the macroalga *Plocamium* spp. (K. Boos, pers. obs.) and attached to drifting surface accumulations of macroalgae (Ashton 2006).

Discussion

Dispersal of *Caprella mutica* within Europe has been rapid and represents ~1,200 km range extension to the west coast of Norway and ~1,000 km to the west coast of Ireland from where it was first recorded in The Netherlands. *C. mutica* has been found on boat hulls, fish cage netting, marina pontoons, navigation buoys, harbour structures and to attached floating macroalgae. Its abundance on artificial structures is notable and may explain its ability to spread rapidly. *C. mutica* lacks a planktonic stage, the young hatch onto the substrate in the form of small adults, and their long-range distribution most probably depends on the movement of floating artificial structures, such as vessels, whilst attachment to floating marine algae may account for more localised movements.

A genetic study indicates two likely routes for the introduction of this species to Europe, including cross-oceanic introductions either directly from Asia or via the Atlantic coasts of the United States (Ashton 2006). Shipping has been identified as an important pathway for the transoceanic introduction of non-native species (Drake and Lodge 2004). Many of the areas in Europe where *C. mutica* has been introduced are close to busy ports suggesting that ballast water transport and/ or hull fouling could be involved. Living *Caprella* spp. have been found in ships' ballast tanks (Carlton 1985) and in sea-chests in a study in New Zealand (Coutts et al. 2003). Within its native environment, *C. mutica* may be found attached to the macroalgae *Ulva* spp. and




Figure 2. European distribution of *Caprella mutica*. Solid circles show confirmed sightings (including dates for first record in the country) and dotted circles show regions where *C. mutica* has not been recorded to date

the filamentous *Cladophora* spp. and these are regularly found attached to ships hulls (Mineur et al. 2007). It has also been seen associated with other algae at high densities on recreational boat hulls (Minchin and Holmes 2006, G Ashton, pers. obs. and R Shucksmith, pers. comm.) and in the Adriatic Sea *Caprella scaura* is thought to have been spread on the hulls of leisure craft (Sconfiatti et al. 2005).

Stock movements of cultured species have also been identified as one means of globally spreading non-native species (Ruiz and Hewitt 2002, Minchin 2007a). Introduction of *C. mutica* to the United States has been potentially linked to the importation of the Pacific Oyster, *C. gigas* for culture purposes (Carlton 1987). Elsewhere there are several examples of species being inadvertently spread with oysters that include algae (Scagel 1956, Druehl 1973, Critchley 1983), molluscs (Cole 1942), bryozoa (De Blauwe and Faasse 2004) and crustaceans (Gruet et al. 1976). The Oosterschelde region in The Netherlands has received extensive shellfish imports and these movements may have been

responsible for the introduction of *C. mutica* to Europe (Wolff 2005). However, this region was regularly sampled between 1990-1995 and *C. mutica* was not found (M. Faasse, pers. obs.). This suggests that the introduction to Europe of *C. mutica* is more likely to be via commercial shipping rather than stock movements of cultured species.

Dispersal of *C. mutica* within Europe has been rapid. It is likely, therefore, that commercial and recreational vessels will continue to spread this species to unoccupied areas within its present range, as well as, to localities beyond where it presently occurs particularly in the summer months, when there is the greatest leisure vessel activity and highest densities of *C. mutica*. On the west coast of Scotland, their abundance can reach 300,000 individuals m^{-2} and fish farm cages (Ashton 2006) and other structures can be dominated by this species between July and August (Cook et al. 2006) (Figure 3). It is possible though, that the rapid range expansion to fish farm installations in Scotland (Willis et al. 2004), Ireland (Tierney et al. 2004) and in Norway (A. Jelmert, pers. comm.) and navigation buoys in Belgium and The Netherlands (F. Kerckhof, pers. obs.) may be with movements of service craft. In California, USA, the appearance of *C. mutica* on oil platforms offshore is thought to be due to transmissions with service craft from harbours (C. Culver, pers. comm.). Such expansion to offshore facilities may create a 'stepping-stone' effect, thereby enhancing opportunities for further range extensions. Spread could also be aided by attachment to drift algae but it is unclear how far *C. mutica* could be dispersed in this way, although algal rafts are known to persist for long durations (Ingolfsson 1995).

The present records of *C. mutica* all occur within the current global range of this species. Its tolerance to a wide range of temperatures and capability of surviving in marine to brackish water (Ashton et al. 2007b) would indicate that it is likely to expand its range further within Europe. It is unlikely to survive in the central and eastern Baltic Sea due to low salinities (FIMR 2006), and based on current knowledge it is not expected to become established in the Mediterranean Sea on account of the high summer seawater temperatures (Cook et al. 2006).

The abundance of *C. mutica* on artificial substrata would indicate that conditions on such structures are in some way particularly suited for



Figure 3. Experimental frame deployed west of Zeebrugge, Belgium in 2007 showing dense aggregations of *Caprella mutica* after 12 weeks immersion (Photo: F. Kerckhof)

their survival and development, such as the lack of benthic predators. Should hull fouling be important in its range expansion it may be expected that their northward expansion in Norway will be slow, when compared with the north and west coasts of France and the Iberian Peninsula where there is a higher level of leisure craft and shipping activity. Other additional factors, that might limit their range expansion and/ or establishment in natural habitats include substrate features (Caine 1978), wave exposure (Takeuchi et al. 1987, Guerra-García 2001), high levels of predation (Guerra-García and García-Gómez 2001) or port toxicity (Ohji et al. 2003).

Conclusions

Caprella mutica has become established in the North Sea, west coasts of Scotland and Ireland, in the Irish Sea and English Channel in less than 14 years. Since this species has a tolerance for a wide range of temperature and salinity, it is likely that it will expand its current range further north in Norway, to several Atlantic islands, the Bay of Biscay and the Iberian Peninsula; but it may not become established in the Mediterranean Sea or in the eastern Baltic Sea, where salinities fall below 19 ppt. *C. mutica* is thought to expand its range rapidly either by attachment to vessels or with drifting algae. It out-competes

native caprellids under laboratory conditions and it is likely to behave in the same way in the field where it attains high densities. Whilst the wider environmental implications of *C. mutica* have not yet been confirmed, it is likely that it has an important impact on benthic communities.

Acknowledgements

We thank all those who provided *Caprella mutica* records: J. Bishop, H. Botnen, R. Fredricksen, J. M. Guerra García, A. Jelmert, E. Leppäkoski, W. Lewis, A. Malej, A. Occhipinti, P. Smith, R. Shucksmith and several members of the Belgian Strandwerkgroep. We also thank Carolynn Culver for information on *C. mutica* distributions in California USA. This research was funded by the Esmée Fairbarin Foundation Marine Aliens Programme, Natural Environment Research Council (NERC) OCEANS 2025, the Scottish Funding Council and EU 6th Framework Programme projects ALARM (contract number GOCE-CT-2003-506675) and DAISIE (contract number SSPI-CT-2003-511202).

References

- Arenas F, Bishop JDD, Carlton JT, Dyrinda PJ, Farnham WF, Gonzalez DJ, Jacobs MW, Lambert C, Lambert G, Nielsen SE, Pederson JA, Porter JS, Ward S and Wood CA (2006) Alien species and other notable records from a rapid assessment survey of marinas on the south coast of England. *Journal of the Marine Biological Association of the United Kingdom* 86: 1329-1337
- Arimoto I (1976) Taxonomic studies of caprellids (Crustacea, Amphipoda, Caprellidae) found in the Japanese adjacent waters. Special publications from the Seto Marine Biological Laboratory Series III, Nippon Printing & Publishing Co., Ltd, Osaka, Japan, 111 pp
- Ashelby CW (2005) The occurrence and distribution of non-native fauna in Harwich Harbour and the Stour and Orwell estuaries, including new records of *Caprella mutica* Schurin 1935 and *Bugula stolonifera* Ryland 1960. *Essex Naturalist* 22: 103-116
- Ashton GV (2006) Distribution and dispersal of the non-native caprellid amphipod, *Caprella mutica* Schurin, 1935. PhD Thesis, Aberdeen, 192 pp
- Ashton GV, Boos K, Shucksmith R and Cook EJ (2006) Rapid assessment of the distribution of marine non-native species in marinas in Scotland. *Aquatic Invasions* 4: 209-213
- Ashton GV, Willis KJ and Cook EJ (2007a) Global Distribution of the Japanese Skeleton Shrimp, *Caprella mutica* (Crustacea, Amphipoda, Caprellidae) with a detailed account of the distribution in Scotland, U.K. *Hydrobiologia* 590: 31-41
- Ashton GV, Willis K, Burrows M and Cook EJ (2007b) Environmental tolerance of *Caprella mutica*: implications for its distribution as a non-native species. *Marine Environmental Research* 64: 305-312
- Buschbaum C and Gutow L (2005) Mass occurrence of an introduced crustacean (*Caprella* cf. *mutica*) in the south-eastern North Sea. *Helgoland Marine Research* 59: 252-253
- Caine EA (1978) Habitat adaptations of North American caprellid Amphipoda (Crustacea). *Biological Bulletin* 155: 288-296
- Carlton JT (1979) History, biogeography, and ecology of the introduced marine and estuarine invertebrates of the Pacific Coast of North America. PhD Thesis, University of California.
- Carlton JT (1985) Transoceanic and interoceanic dispersal of coastal marine organisms: the biology of ballast water. *Oceanogr Mar Biol. An Annual Review* 23: 313-371
- Carlton JT (1987) Patterns of transoceanic marine biological invasions in the Pacific ocean. *Bulletin Marine Science* 41: 452-465
- Carlton JT (1996) Biological Invasions and Cryptogenic Species. *Ecology* 77: 1653-1655
- Cohen AN and Carlton JT (1995) Nonindigenous aquatic species in an United States Estuary: A case study of the biological invasions of the San Francisco Bay and Delta. A Report for the United States fish and wildlife service, Washington D.C. and The national sea grant college program Connecticut sea grant. Available online: <http://www.anstaskforce.gov/sfinvade.htm>, 246 pp
- Cole HA (1942) The American whelk tingle, *Urosalpinx cinerea* (Say), on British oyster beds. *Journal of the Marine Biological Association of the United Kingdom* 25: 477-508
- Cook EJ, Black KD, Sayer MDJ, Cromey CJ, Angel DL, Spanier E, Tsemel A, Katz T, Eden N, Karakassis I, YTsapakis M, Apostolaki ET and Malej A (2006) The influence of caged mariculture on the early development of sublittoral fouling communities: a pan-European study. *ICES Journal of Marine Science* 63: 637-649
- Cook EJ, Willis KJ and Lozano-Fernandez M (2007) Survivorship, Growth and Reproduction of the Non-Native *Caprella mutica* Schurin (Crustacea: Amphipoda). *Hydrobiologia* 590: 55-64
- Coutts ADM, Moore KM and Hewitt CL (2003) Ships' sea-chests: an overlooked transfer mechanism for non-indigenous marine species? *Marine Pollution Bulletin* 46: 1504-1515
- Critchley AT (1983) *Sargassum muticum*: a taxonomic history including world-wide and western Pacific distributions. *Journal of the Marine Biological Association of the United Kingdom* 63: 617-625
- De Blauwe H and Faasse MA (2004) *Smittoidea prolifica* Osburn, 1952 (Bryozoa, Cheilostomatida), a Pacific bryozoan introduced to The Netherlands (Northeast Atlantic). *Bulletin van het Koninklijk Belgisch Instituut voor Natuurwetenschappen, Biologie* 74: 33-39
- Drake JM and Lodge DM (2004) Global hot spots of biological invasions: evaluating options for ballast-water management. *Proceedings of the Royal Society London B* 271: 575-580
- Druel L (1973) Marine transplantations. *Science* 179: 12
- FIMR (2006) FIMR monitoring of the Baltic Sea environment - Annual Report 2006. Finnish Institute for Marine Research, Helsinki
- Gruet Y, Heral M and Robert J-M (1976) Premières observations sur l'introduction de la faune associée au naissain d'huîtres Japonaises *Crassostrea gigas* (Thunberg), importé sur la côte Atlantique Française. *Cahiers de Biologie Marine* 17: 173-184

- Guerra-García JM (2001) Habitat use of the Caprellidea (Crustacea : Amphipoda) from Ceuta, North Africa. *Ophelia* 55: 27-38
- Guerra-García JM and García-Gómez J (2001) The spatial distribution of Caprellidae (Crustacea: Amphipoda): a stress bioindicator in Ceuta (North Africa, Gibraltar Area). *Marine Ecology- Pubblicazioni Della Stazione Zoologica Di Napoli* 22: 357-367
- Heilscher S (2000) Sustainability of the brown seaweed *Laminaria saccharina* for cultivation in the effluents of an Atlantic salmon ongrowth farm. A field study in Southeastern Norway, Institut für Biologie, Freiburg im Breisgau
- Héral M (1990) Traditional oyster culture in France. In: Barnabé (ed) *Aquaculture*, pp 342-387, Ellis-Horwood, London
- Ingólfsson A (1995) Floating clumps of seaweed around Iceland - natural microcosms and a means of dispersal for shore fauna. *Marine Biology* 122: 13-21
- Kawashima H, Takeuchi I and Ohnishi M (1999) Fatty acid compositions in four of caprellid amphipod species (Crustacea) from Otsuchi and Mutsu bays in northern Japan. *Journal of Japanese Oil Chemistry Society* 48: 595-599
- Kerckhof F, Haelters J and Gollasch S (2007) Alien species in the marine and brackish ecosystem: the situation in Belgian waters. *Aquatic Invasions* 2(3): 243-257
- Loosanoff VL (1975) Introduction of *Codium* to New England waters. *Fisheries Bulletin* 73: 215-218
- Marelli DC (1981) New records for Caprellidae in California, and notes on a morphological variant of *Caprella verrucosa* Boeck, 1871. *Proceedings of the Biological Society Washington* 94: 654-662
- Minchin D (2007a) Aquaculture and transport in a changing environment: Overlap and links in the spread of alien biota. *Marine Pollution Bulletin* 55: 302-313
- Minchin D (2007b) Rapid coastal survey for targeted alien species associated with floating pontoons in Ireland. *Aquatic Invasions* 2: 63-70
- Minchin D and Holmes JMC (2006) The first record of *Caprella mutica* Schurin 1935 (Crustacea: Amphipoda) from the east Coast of Ireland. *Irish Naturalists' Journal* 28: 321-323
- Mineur F, Belsher T, Johnson MP, Maggs CA and Verlaque M (2007) Experimental assessment of oyster transfers as a vector for macroalgal introductions. *Biological Conservation* 137: 237-247
- Nishimura S (1995) Guide to the Seashore Animals of Japan with Colour Pictures and Keys. Vol. II. Hoikusha, Japan
- O'Reilly M (2007) The Japanese Macho Skeleton Shrimp (*Caprella mutica*) in the Clyde Estuary. *Glasgow Naturalist* 24: 156
- Ohji M, Arai T and Miyazaki N (2003) Chronic effects of tributyltin on the caprellid amphipod *Caprella danilevskii*. *Marine Pollution Bulletin* 46: 1263-1272
- Platvoet D, de Bruyne RH and Gmelig Meyling AW (1995) Description of a new *Caprella*-species from the Netherlands: *Caprella macho* nov.spec. (Crustacea, Amphipoda, Caprellidae). *Bulletin of the Zoological Museum, University of Amsterdam* 15: 1-4
- Rodríguez G and Suárez H (2001) Anthropogenic dispersal of decapod crustaceans in aquatic environments. *Interciencia* 26 282
- Ruiz GM and Hewitt CL (2002) Toward understanding patterns of coastal marine invasions: a prospectus. In: Leppäkoski E, Gollasch S and Olenin S (eds) *Invasive aquatic species of Europe. Distribution, impacts and management*, Kluwer Academic Publishers, Dordrecht, The Netherlands
- Scagel RF (1956) Introduction of a Japanese alga, *Sargassum muticum* into the northeast Pacific. *Fisheries Research Papers. Department of Fisheries, State of Washington* 1: 49-59
- Schrey I (2006) Verbreitung des eingeschleppten Gespensterkrebses *Caprella mutica* in der Deutschen Bucht und seine potentielle Ansiedlung in natürlichen Habitaten. Diploma Thesis, Rostock University
- Schurin A (1935) Zur Fauna der Caprelliden der Bucht Peter der Grossen (Japanisches Meer). *Zoologischer Anzeiger* 122: 198-203
- Sconfiatti R, Mangili F, Savini D and Occhipinti-Ambrogi A (2005) Diffusion of the alien species *Caprella scaura* Templeton, 1836 (Amphipoda:Caprellidae) in the Northern Adriatic Sea. *Biologia Marina Mediterranea* 12: 335-337
- Shucksmith R (in press) Biological Invasions: The role of biodiversity in determining community susceptibility to invasion PhD Thesis, Aberdeen
- Stoner D, Ben-Shlomo R, Rinkevich B and Weissman I (2002) Genetic variability of *Botryllus schlosseri* invasions to the east and west coasts of the USA. *Marine Ecology Progress Series* 243: 93-100
- Takeuchi I, Kuwabara R, Hirano R and Yamakawa H (1987) Species compositions of the Caprellidae (Crustacea: Amphipoda) of the *Sargassum* zone on the Pacific coast of Japan. *Bulletin of Marine Science* 41: 253-267
- Tierney TD, Kane F, Naughton O, Kennedy S, O'Donohoe P, Copley L and Jackson D (2004) On the occurrence of the caprellid amphipod, *Caprella mutica* Schurin 1935, in Ireland. *Irish Naturalists' Journal* 27: 437-439
- Vassilenko SV (1967) Fauna of Caprellidae (Amphipoda) of the Possjet Bay (Sea of Japan) and some data on their ecology. *Issledovanija Fauny Morei (Explorations of the fauna of the seas of the USSR)*, Biotzenozy Zalika Possjet Japanskovo Morja 5: 196-229
- WGITMO (2003) Report of the ICES Working Group on Introductions and Transfers of Marine Organisms, Vancouver, Canada, 158 pp
- Willis KJ, Cook EJ, Lozano-Fernandez M and Takeuchi I (2004) First record of the alien caprellid amphipod, *Caprella mutica*, for the UK. *Journal of the Marine Biological Association of the United Kingdom* 84: 1027-1028
- Wolff WJ (2005) Non-indigenous marine and estuarine species in The Netherlands. *Zoologische Mededelingen Leiden* 79: 1-116

European expansion of *Caprella mutica*

Annex

European distribution records of *Caprella mutica* including location, latitude and longitude, date of reporting and site description (if known). Confirmed non-sightings are also included

Location/Country	Geographic coordinates		Date	Site description	Source
	Latitude	Longitude			
Neeltje Jans Binnenhaven, Netherlands	51°37' N	03°41' E	Summer, 1994	Harbour	M. Faasse (pers. obs.)
Burghsluis, Netherlands	51°40' N	03°45' E	1995	Pontoon & scaffolding	Platvoet et al. 1995
Neeltje Jans Buitenhaven, Netherlands	51°37' N	03°41' E	April, 1996	Harbour	M. Faasse (pers. obs.)
Goesse Sas, Netherlands	51°33' N	03°56' E	June, 1996	Seadyke	M. Faasse (pers. obs.)
Katse Hoek, Netherlands	51°34' N	03°54' E	June, 1996	Seadyke	M. Faasse (pers. obs.)
's Gravenhoek, Netherlands	51°39' N	03°48' E	July, 1996	Seadyke	M. Faasse (pers. obs.)
Kattendijke, Netherlands	51°32' N	03°58' E	July, 1996	Seadyke	M. Faasse (pers. obs.)
Borssele, Netherlands	51°25' N	03°43' E	August, 1997	Power station	M. Faasse (pers. obs.)
Zierikzee west, Netherlands	51°38' N	03°53' E	September, 1997	Seadyke	M. Faasse (pers. obs.)
Off Zeebrugge east, Belgium	51°37' N	3°41' E	January, 1998	Open water buoy (S12)	F. Kerckhof (pers. obs.)
Off Zeebrugge west, Belgium	51°2' N	3°00' E	March, 1998	Open water buoy (SWW)	F. Kerckhof (pers. obs.)
Off Zeebrugge west, Belgium	51°23' N	2°58' E	November, 1998	Open water buoy (S2)	F. Kerckhof (pers. obs.)
Off Zeebrugge east, entrance river Scheldt, Belgium	51°23' N	3° 18' E	July, 1998	Open water buoys (W1 and SW)	F. Kerckhof (pers. obs.)
Port of Zeebrugge, Belgium	51°19' N	3°12' E	July, March, November, 1998	Marina and buoys (Z10, Z8, ZoK N)	F. Kerckhof (pers. obs.) H. De Blauwe (pers. comm.)
Neeltje Jans buoy, Netherlands	51°37' N	03°42' E	April, 1998	Open water buoy	M. Faasse (pers. obs.)
Neeltje Jans Vluchthaven, Netherlands	51°38' N	03°42' E	May, 1998	Aquaculture	M. Faasse (pers. obs.)
Zeebrugge, Belgium	51°19' N	03°12' E	1999	Marina	H. De Blauwe (pers. comm.)
Off Zeebrugge in front of harbour entrance, Belgium	51°22' N	03°10' E	May, 1999	Open water buoy (WZ)	F. Kerckhof (pers. obs.)
Off Zeebrugge, northwest, Belgium	51°24' N	03°03' E	July, 1999	Open water buoy (S3)	F. Kerckhof (pers. obs.)
Hordaland, Norway	60°14' N	06°00' E	1999	Harbour	Heilscher 2000
Clyde Sea, Scotland, UK	55°57' N	04°52' W	August, 1999	Navigation buoy	O'Reilly 2007
Port of Zeebrugge, Belgium	51°19' N	03°12' E	January, October, 2000	Port (ZA1, ZA4)	F. Kerckhof (pers. obs.)
Goesse Meer, Netherlands	51°31' N	03°55' E	July, 2000	Inland water	M. Faasse (pers. obs.)
Breskens, Netherlands	51°24' N	03°34' E	October, 2000	Marina	M. Faasse (pers. obs.)
Bruinisse, Netherlands	51°40' N	04°06' E	November, 2000	Harbour	M. Faasse (pers. obs.)
List, Sylt, Germany	55° 00' N	08°26' E	2000	Unknown	Buschbaum and Gutow 2005
Helgoland, Germany	54°10' N	07°53' E	2000	Unknown	Buschbaum and Gutow 2005
Oban, Scotland, UK	56°26' N	05°26' W	July, 2000	Fish farm	Willis et al. 2004

Annex (continued)

Location/Country	Geographic coordinates		Date	Site description	Source
	Latitude	Longitude			
Wissenkerke east, Netherlands	51°36' N	03°47' E	August, 2001	Seadyke	M. Faasse (pers. obs.)
Goesse Sas Marina, Netherlands	51°32' N	03°56' E	November, 2001	Marina	M. Faasse (pers. obs.)
Wissenkerke, Netherlands	51°36' N	03°45' E	June, 2001	Seadyke	M. Faasse (pers. obs.)
Zoetersbout, Netherlands	51°39' N	04°06' E	June, 2001	Seadyke	M. Faasse (pers. obs.)
Oostende, Belgium	51°13' N	02°54' E	April, 2002	Small buoy in front of harbour	F. Kerckhof (pers. obs.)
Off Zeebrugge east, Belgium	51°24' N	03°18' E	April, 2002	Open water buoy (S12)	F. Kerckhof (pers. obs.)
Harwich, England, UK	51°56' N	01°16' E	June, 2002	Harbour	Ashelby 2005
Vlissingen, Netherlands	51°26' N	03°35' E	July, 2002	Marina	M. Faasse (pers. obs.)
Zierikzee, Netherlands	51°38' N	03°55' E	July, 2002	Seadyke	M. Faasse (pers. obs.)
Wemeldinge, Netherlands	51°31' N	04°00' E	August, 2002	Marina	M. Faasse (pers. obs.)
Alesund, Norway	62°29' N	06°23' E	2002		WGITMO 2003, A. Jelmert (pers. comm.)
Off Zeebrugge in front of harbour entrance, Belgium	51°22' N	03°10' E	August, 2003	Open water buoy (WZ) Small	F. Kerckhof (pers. obs.)
Blankenberge, Belgium	51°19' N	03°07' E	October, 2003	temporary buoys in front of harbour	F. Kerckhof (pers. obs.)
Port of Zeebrugge, Belgium	51°19' N	03°12' E	July, November, 2003	Port (buoys ZoK N, ZW1)	F. Kerckhof (pers. obs.)
Southampton, England, UK	50°53' N	01°23' W	2003	Harbour	L. Baldock & M. Marley (pers. comm.)
Bertraghboy Bay, Ireland	53°29' N	10°04' W	2003	Fish farm	Tierney et al. 2004
Hidra, Norway	58°15' N	06°32' E	2003	Aquaculture	A. Jelmert (pers. comm.)
Austevoll, Norway	60°06' N	05°15' E	2003	Aquaculture	A. Jelmert (pers. comm.)
Shetland Is, Scotland, UK	60°22' N	01°12' W	2003	Mussel lines	G. Duncan (pers. comm.)
Anglesey, Wales, UK	53°18' N	04°37' W	August, 2003	Mooring line	T. Stoker (pers. comm.)
Off Zeebrugge, west, Belgium	51°02' N	03°02' E	September, 2004	Open water buoy (WBN)	F. Kerckhof (pers. obs.)
Southampton, England, UK	50°53' N	01°23' W	2004	Pontoons	Arenas et al. 2006
Poole Quay, England, UK	50°42' N	01°59' W	2004	Pontoons	Arenas et al. 2006
Le Havre, France	49°29' N,	00°07' E	2004	Harbour	G. Breton (pers. comm.)
List, Sylt, Germany	55°00' N	08°26' E	October, 2004	Pontoons	Buschbaum and Gutow 2005
Helgoland, Germany	54°10' N	07°53' E	October, 2004	Pontoons	Buschbaum and Gutow 2005
Dunstaffnage Bay, Scotland, UK	56°27' N	05°26' W	April, 2004	Marina	Ashton 2006
Lamlash, Scotland, UK	55°31' N	05°07' W	May, 2004	Fish Farm	Ashton 2006
Clyde Basin, Scotland, UK	55°57' N	04°52' W	May, 2004	Fish Farm	Ashton 2006
Loch Fyne, Scotland, UK	55°56' N	05°23' W	May, 2004	Fish Farm	Ashton 2006
Fishnish, Scotland, UK	56°30' N	05°48' W	May, 2004	Fish Farm	Ashton 2006

European expansion of *Caprella mutica*

Annex (continued)

Location/Country	Geographic coordinates		Date	Site description	Source
	Latitude	Longitude			
Lismore, Scotland, UK	56°30' N	05°30' W	May, 2004	Fish Farm	Ashton 2006
Shuna, Scotland, UK	56°12' N	05°36' W	May, 2004	Fish Farm	Ashton 2006
Kinlochleven, Scotland, UK	56°42' N	04°58' W	May, 2004	Fish Farm	Ashton 2006
Loch Sunart, Scotland, UK	56°41' N	05°46' W	May, 2004	Fish Farm	Ashton 2006
Lochboisdale, Scotland, UK	57°09' N	07°18' W	May, 2004	Fish Farm	Ashton 2006
Loch Carnan, Scotland, UK	57°22' N	07°16' W	May, 2004	Fish Farm	Ashton 2006
Loch Kishorn, Scotland, UK	57°21' N	05°40' W	May, 2004	Fish Farm	Ashton 2006
Loch Torridon, Scotland, UK	57°34' N	05°46' W	May, 2004	Fish Farm	Ashton 2006
Loch Maddy, Scotland, UK	57°36' N	07°09' W	May, 2004	Fish Farm	Ashton 2006
Cheese Bay, Scotland, UK	57°39' N	07°06' W	May, 2004	Fish Farm	Ashton 2006
Loch Ewe, Scotland, UK	57°48' N	05°37' W	May, 2004	Fish Farm	Ashton 2006
Loch Tarbert, Scotland, UK	55°51' N	05°26' W	May, 2004	Fish Farm	Ashton 2006
Loch Seaforth, Scotland, UK	57°57' N	06°43' W	May, 2004	Fish Farm	Ashton 2006
Loch Shell, Scotland, UK	58°00' N	06°30' W	May, 2004	Fish Farm	Ashton 2006
Loch Erisort, Scotland, UK	58°05' N	06°29' W	May, 2004	Fish Farm	Ashton 2006
Loch Roag, Scotland, UK	57°26' N	06°35' W	May, 2004	Fish Farm	Ashton 2006
Calbha Bay, Scotland, UK	58°17' N	05°07' W	May, 2004	Fish Farm	Ashton 2006
Loch Laxford, Scotland, UK	58°22' N	05°01' W	May, 2004	Fish Farm	Ashton 2006
Off Zeebrugge west, Belgium	51°23' N	02°46' E	June, 2005	Open water buoy (VG)	F. Kerckhof (pers. obs.)
Off Zeebrugge, northwest, Belgium	51°24' N	03°03' E	September, 2005	Open water buoy (S3)	F. Kerckhof (pers. obs.)
Off Zeebrugge west, Belgium	51°22' N	03°05' E	September, 2005	Open water buoy (Nippon)	F. Kerckhof (pers. obs.)
Off Oostende harbour, Belgium	51°14' N	02°53' E	October, 2005	Open water buoy (Buitenstroombank)	F. Kerckhof (pers. obs.)
Horns Rev Windfarm, Denmark	55°27' N	08°26' E	2005	Windfarm	R. Frederiksen (pers. comm.)
Wangerooge, Germany	53°47' N	07°54' E	July, 2005	Pontoons	Schrey 2006
Hoernum, Sylt, Germany	54°45' N	08°17' E	October, 2005	Pontoons	Schrey 2006
Dun Laoghaire, Dublin Bay, Ireland	54°03' N	06°11' W	2005	Marina pontoons & floating algae	Minchin and Holmes 2006
Anna Jacobapolder, Netherlands	51°39' N	04°06' E	July, 2005	Harbour	M. Faasse (pers. obs.)
Wolphaartsdijk, Netherlands	51°33' N	03°49' E	July, 2005	Seadyke	M. Faasse (pers. obs.)
Texel, Netherlands	53°02' N	04°41' E	March, 2005	Buoy (MG 10)	F. Kerckhof (pers. obs.)
Texel, Netherlands	53°02' N	04°41' E	October, 2005	Buoy (MG 1A)	F. Kerckhof (pers. obs.)

Annex (continued)

Location/Country	Geographic coordinates		Date	Site description	Source
	Latitude	Longitude			
Marsdiep, Netherlands	53°04' N	05°28' E	2005	Mussel Lines	W. Lewis (pers. comm.)
Malzwin, Netherlands	53°03' N	05°27' E	2005	Mussel Lines	W. Lewis (pers. comm.)
Loch Linnhe, Scotland, UK	56°31' N	05°26' W	May, 2005	Floating macroalgae	Ashton 2006
Loch Beag, Scotland, UK	56°53' N	05°44' W	June, 2005	Mussel Lines	E. Cook (pers. obs.)
Loch Creran, Scotland, UK	56°31' N	05°20' W	August, 2005	Boat hull	R. Shucksmith (pers. comm.)
Oban, Scotland, UK	56°25' N	05°30' W	August, 2005	Mooring lines, pontoons, kelp beds	R. Shucksmith (pers. comm.)
Off Zeebrugge, east, Belgium	51°24' N	03°18' E	March, 2006	Open water buoys (S12 and S10)	F. Kerckhof (pers. obs.)
Off Zeebrugge, northwest, Belgium	51°23' N	03°05' E	July, 2006	Open water buoy (S5)	F. Kerckhof (pers. obs.)
Port of Zeebrugge, Belgium	51°19' N	03°12' E	October, 2006	Port buoy (ZA2)	F. Kerckhof (pers. obs.)
Off Nieuwpoort, Belgium	51°13' N	02°38' E	November, 2006	Open water buoy from mussel cultivation project	F. Kerckhof (pers. obs.)
Off Zeebrugge, east, Shipping route, Belgium	51°24' N	02°44' E	November, 2006	Open water buoy (GZ)	F. Kerckhof (pers. obs.)
Off Zeebrugge, west, Belgium	51°22' N	03°07' E	December, 2006	Open water buoy (A2)	F. Kerckhof (pers. obs.)
Cork Harbour, Ireland	51°50' N	08°16' W	July, August, 2006	Marina pontoons	Minchin 2007b
Carlingford Lough, Ireland	54°03' N	06°11' W	July, August, 2006	Marina pontoons	Minchin 2007b
Ardrossan, Scotland, UK	55°38' N	04°49' W	August, 2006	Marina pontoons	Ashton et al. 2006
Troon, Scotland, UK	55°33' N	04°42' W	August, 2006	Marina pontoons	Ashton et al. 2006
Peterhead, Scotland, UK	57°29' N	01°47' W	August, 2006	Marina pontoons	Ashton et al. 2006
Port Edgar, Scotland, UK	55°60' N	03°25' W	August, 2006	Marina pontoons	Ashton et al. 2006
Lossiemouth, Scotland, UK	57°43' N	03°16' W	August, 2006	Marina pontoons	Ashton et al. 2006
Ardfern, Scotland, UK	56°11' N	05°32' W	August, 2006	Marina pontoons	Ashton et al. 2006
Croabh Haven, Scotland, UK	56°13' N	05°33' W	August, 2006	Marina pontoons	Ashton et al. 2006
Off Zeebrugge west, Belgium	51°32' N	02°22' E	March, 2007	Open water buoy (VG2)	F. Kerckhof (pers. obs.)
Off Zeebrugge, Belgium	51°27' N	02°09' E	July, 2007	Open water buoy (NE Akkaert)	F. Kerckhof (pers. obs.)

European expansion of *Caprella mutica*

Annex (continued)

Location/Country	Geographic coordinates		Date	Site description	Source
	Latitude	Longitude			
Koksijde, Belgium	51°07' N	02°36' E	October, 2007	Small temporary buoy	F. Kerckhof (pers. obs.)
Zeebrugge West, Belgium	51°22' N	03°07' E	October, 2007	Experimental Frame	F. Kerckhof (pers. obs.)
Plymouth, England, UK	50°22' N	04°08' W	July, 2007	Marina pontoons	J. Bishop (pers. comm.)
Helgoland, Germany	54°10' N	07°53' E	October, 2007	Macroalga, Plocamium spp.	K. Boos (pers. obs.)
Sture, Norway	60°49' N	03°30' E	2007	Abandoned fish farm	H. Botnen (pers. obs.)
Nyhamna, Norway	62°50' N	06°54' E	2007	New harbour for gas processing plant	H. Botnen (pers. obs.)
Loch Fyne, Scotland, UK	56°09' N	05°11' W	May, 2007	Fishing boat hull	P. Smith (pers. comm.)
Loch Melfort, Scotland, UK	56°15' N	05°32' W	July, 2007	Boat hull	E. Cook (pers. obs.)
Largs, Scotland, UK	55°46' N	04°51' W	August, 2007	Marina pontoons	E. Cook (pers. obs.)
Regions where <i>C. mutica</i> has not been found to date					
HELCOM Baltic, Finland					E. Leppäkoski (pers. comm.)
Dunkerque, Calais, Boulogne and Boullonnais coast, France				Ports	Belgium Strandwerkgroep (pers. comm.)
Granville, Normandy, France				Port	Belgium Strandwerkgroep (pers. comm.)
Brittany, France				Coastal survey	Belgium Strandwerkgroep (pers. comm.)
Venice Lagoon, Italy	45°27' N	12°19' E			A. Occhipinti (pers. comm.)
Cadiz, Spain	36°31' N	6°17' W			J. M. Guerra García (pers. comm.)
Alicante, Spain	38°25' N	0°22' W			J. M. Guerra García (pers. comm.)
Adriatic Sea, Slovenia/Croatia					A. Malej (pers. comm.)
Basque coast, Spain					Belgium Strandwerkgroep (pers. comm.)
La Coruna and adjacent region, Spain	43°22' N	8°23' W			F. Kerckhof (pers. obs.)
Ria de Vigo and adjacent region, Spain	42°13' N	8°46' W			F. Kerckhof (pers. obs.)